arXiv:1509.07595v1 [math.AP] 25 Sep 2015

UNIQUENESS OF POSITIVE SOLUTIONS OF A n-LAPLACE EQUATION IN A BALL IN $\mathbb{R}^n$ WITH EXPONENTIAL NONLINEARITY

ADIMURTHI, KARTHIK A, AND JACQUES GIACOMONI

Abstract. Let $n \geq 2$ and $\Omega \subset \mathbb{R}^n$ be a bounded domain. Then by Trudinger-Moser embedding, $W_0^{1,n}(\Omega)$ is embedded in an Orlicz space consisting of exponential functions. Consider the corresponding semi linear n-Laplace equation with critical or sub-critical exponential nonlinearity in a ball $B(R)$ with dirichlet boundary condition. In this paper, we prove that under suitable growth conditions on the nonlinearity, there exists an $\gamma_0 > 0$, and a corresponding $R_0(\gamma_0) > 0$ such that for all $0 < R < R_0$, the problem admits a unique non degenerate positive radial solution u with $\|u\|_\infty \geq \gamma_0$.

1. Introduction

Let $B(R) \subset \mathbb{R}^n$ be the ball of radius R with center at the origin and $1 < p \leq n$ with $f \in C^0[0, +\infty)$. Consider the following problem

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p-2}\nabla u) = f(u) & \text{in } B(R) \\ u = 0 \quad \text{on } \partial B(1), \ u > 0 & \text{in } B(R). \end{cases} \tag{1.1}$$

Existence of a solution to Problem (1.1) has been studied extensively under the banner of Emden-Fowler type equations or Yamabe type equations. In this paper we make some progress regarding the question of uniqueness.

History for $p = 2$. Let $\lambda > 0$ and define

$$f(u) = \begin{cases} u^r + \lambda u & \text{if } n \geq 2 \text{ and } 1 < r < \infty \\ \lambda\, h(u)e^{u^q} & \text{if } n = 2 \text{ and } 1 \leq q \leq 2, \end{cases} \tag{1.2}$$

where $h \in C^1(0, \infty)$ with $h(0) = 0$ and is of lower order growth than e^{u^q}. In this context, H. Brezis raised the following question:

"Does (1.1) *admit utmost one solution?"*

From Gidas-Ni-Nirenberg [10], we know that any solution of (1.1) with $p = 2$ is radial. Using this and by scaling argument when $\lambda = 0$, in [10], it was proven that (1.1) with $p = 2$ admits utmost one solution. When $\lambda \neq 0$, their approach fails to yield uniqueness. Using the Pohozaev's identity for $n \geq 2$ and $r < \infty$, the uniqueness question was answered in [5, 9, 12, 16, 15]. In [9, 17], the authors used the Pohozaev's identity in an ingenious way to obtain uniqueness for a wide class of nonlinearities $f(u)$.

Next, consider the case when $n = 2$ and the nonlinearity is of exponential type. If $f(u) = ue^u$, then uniqueness was proved in [2, 17]. In [17], they again made use of Pohozaev's identity, though unfortunately, their technique cannot be extended if $q > 1$. Hence the question remained:

"What happens if $n = 2$ and $1 < q \leq 2$?"

In this paper we try to answer the uniqueness by adopting the linearization technique from [2] and the asymptotic analysis of Atkinson-Peletier [7, 11] (see also Volkmer [18]). Under suitable conditions on f we show that there exists a $\lambda_0 > 0$ such that for $p = n$, $0 < \lambda < \lambda_0$, $q \in (1, \frac{n}{n-1}]$ and $\|u\|_\infty >$ 'some large quantity', the radial solutions of (1.1) are unique and nondegenerate. For the existence and nonexistence of positive solutions see [1, 3].

Hypothesis 1. Let $f \in C[0,\infty) \cap C^3(0,\infty)$ with $f(u) > 0$ for $u > 0$. Let $a > 0$, $\lambda > 0$, $\rho \in C^3(0,\infty)$ and $1 < q \leq \dfrac{n}{n-1}$ be such that

$$f(u) = \lambda e^{g(u)}, \quad g(u) = au^q + \rho(u). \tag{1.3}$$

Let $\rho^{(k)}$ denote the k-th derivative of ρ. Assume that there exist $0 < \alpha \leq 1$, $\beta > 0$ such that

$$\begin{aligned} f(s) - f(0) &= \mathcal{O}\left(s^{\beta}\right) \text{ as } s \to 0^+ \\ f'(s) &= \mathcal{O}\left(s^{-1+\alpha}\right) \text{ as } s \to 0^+. \end{aligned} \tag{1.4}$$

Now the following assumptions will be stated as and when they are used in the theorems:

(H1) $\lim\limits_{\gamma\to\infty} \dfrac{\rho^{(k)}(\gamma)}{\gamma^{q-k}} = 0$ for $k \in \{0,1,2,3\}$.

(H2) There exists a $b \in \mathbb{R}$ such that

$$\varliminf_{\gamma\to\infty} g(\gamma) - \left(\frac{n-1}{n}\right)\gamma g'(\gamma) \geq b$$

(H3) $\lim\limits_{\gamma\to\infty} \dfrac{g'(\gamma)}{g''(\gamma)\left[\log g'(\gamma)\right]^4}\left(g'(\gamma) - (n-1)\gamma g''(\gamma)\right) = \infty.$

Here g', g'' denote the first and second derivatives of g.

We now have the following main result:

Theorem 1.1. *Assume f satisfies (H1) and (H3) from Hypothesis 1 and that there exists a $p \geq 0$ and a constant $C_p \neq 0$ such that*

$$\lim_{u\to 0^+} \frac{f(u)}{u^p} = C_p. \tag{1.5}$$

Then there exists a $\lambda_0 > 0$, $\gamma_0 > 0$ such that for all $\lambda \in (0,\lambda_0)$

$$\begin{cases} -\operatorname{div}(|\nabla u|^{n-2}\nabla u) = \lambda f(u) & \text{in } B(1), \\ u > 0, \quad u \text{ radial in } B(1), \\ u = 0 \quad \text{on } \partial B(1), \end{cases} \tag{1.6}$$

admits a unique solution satisfying $\|u\|_\infty > \gamma_0$. Furthermore, the solution is non degenerate.

Remark 1.2. *By integration, (H1) and (H3) of Hypothesis 1 implies (H2). In fact, one can find $\gamma_0 > 0$, $M > 0$ and a constant $c \in \mathbb{R}$ such that for all $\gamma \geq \gamma_0$, we have the estimate*

$$g(\gamma) - \left(\frac{n-1}{n}\right)\gamma g'(\gamma) \geq M(\log\gamma)^5 + c.$$

It is easy to see in the subcritical case, that is when $1 < q < \dfrac{n}{n-1}$, (H3) of Hypothesis 1 always holds. The question now remains if one can remove the hypothesis (H3)!

First consequence of Theorem 1.1 is that large solution coincides with the solution obtained by the moutainpass lemma and we have the following Corollary:

Corollary 1.3. *Assume that the hypothesis in Theorem 1.1 are satisfied. Then there exists $\lambda_0 > 0$ and $\gamma_0 > 0$ such that for all $\lambda \in (0,\lambda_0)$, solutions to (1.6) satisfying $\|u\|_\infty > \gamma_0$ has a Morse index equal to one.*

Examples. We give some examples of f that satisfy conditions in Hypothesis 1.

(i) Let $f(u) = u^p e^{u^q}$, $0 \le p < \infty$, then we have

$$g(\gamma) = \gamma^q + p\log(\gamma), \quad \rho(\gamma) = p\log(\gamma),$$

$$g(\gamma) - \left(\frac{n-1}{n}\right)\gamma g'(\gamma) = \left(\frac{n-1}{n}\right)\gamma^q\left(\frac{n}{n-1} - q\right) + p\log(\gamma) - \frac{p(n-1)}{n}$$

$$g'(\gamma) - (n-1)\gamma g''(\gamma) = (n-1)q\gamma^{q-1}\left(\frac{n}{n-1} - q\right) + \frac{pn}{\gamma}.$$

Therefore, if $1 < q \le \dfrac{n}{n-1}$, we see that (H1) and (H2) of Hypothesis 1 is satisfied. If $1 < q < \dfrac{n}{n-1}$, then (H1), (H2) and (H3) are satisfied, but when $q = \dfrac{n}{n-1}$, (H3) is not satisfied.

(ii) Let $f(u) = \lambda u^p e^{u^q + \beta u}$ with $\beta > 0$. In this case, (H1), (H2) and (H3) are satisfied for $1 \le q \le \dfrac{n}{n-1}$.

From the Picone's Identity [6], assuming (1.5), there exist a $\lambda_1 > 0$, $\gamma_1 > 0$ such that (see Lemma 2.1):

- If $0 \le p < n-1$, then for $\lambda \in (0, \lambda_1)$, there exists a solution u of (1.6) with $\|u\|_\infty < \gamma_1$.
- If $p \ge n-1$, then for $\lambda \in (0, \lambda_1)$, there does not exist a solution with $\|u\|_\infty < \gamma_1$.

Hence by plotting the solution curve $\|u_\lambda\|_\infty$ with respect to λ, we have the following diagrams.



Case $0 \le p < n-1$.



Case $p = n-1$.



Case $p > n-1$.

Remark 1.4. *We now recall the Trudinger-Moser imbedding* [13]*: let $\Omega \in \mathbb{R}^n$ be a bounded open set, then the Trudinger-Moser imbedding says*

$$A = \sup_{\substack{u \in W_0^{1,n}(\Omega) \\ \|u\| \le 1}} \int_\Omega e^{(n\omega_n^{\frac{1}{n-1}})|u|^{\frac{n}{n-1}}} < \infty,$$

where $\omega_n =$ *volume of* S^{n-1} *and* $n\omega_n^{\frac{1}{n-1}}$ *is the best constant. It was shown by* CARLESON-CHANG [8] *that if* Ω *is a ball, then* A *is achieved by a radial function* $u_0 \in W_0^{1,n}(\Omega)$ *with* $u_0 > 0$ *satisfying the equation*

$$\left\{\begin{array}{ll} -\operatorname{div}(|\nabla u_0|^{n-2}\nabla u_0) = \lambda_0 u_0^{\frac{1}{n-1}} e^{(n\omega^{\frac{1}{n-1}})u_0^{\frac{n}{n-1}}} & \textit{in } \Omega, \\ u_0 > 0, \quad u_0 \textit{ radial in } \Omega, & \\ u_0 = 0 \quad \textit{on } \partial\Omega, & \end{array}\right. \tag{1.7}$$

for some $\lambda_0 > 0$.

Conjecture. Maximizers are unique!

In case of $n = 2$, *if one can show that* (1.6) *admits utmost one solution for all* $\lambda > 0$, *then the above conjecture is positively answered.*

Remark 1.5. *In* [4], *the following singular Trudinger Moser Imbedding has been shown:*

$$\sup_{\|u\|_{W_0^{1,p}(\Omega)} \leq 1} \int_\Omega \frac{e^{\alpha|u|^{\frac{n}{n-1}}}}{|x|^\beta} < \infty,$$

holds if and only if α *and* β *satisfy* $\dfrac{\alpha}{\delta_n} + \dfrac{\beta}{n} \leq 1$ *where* $\delta_n := n\omega_n^{\frac{1}{n-1}}$.

In view of this, consider the following singular semilinear equation

$$\left\{\begin{array}{ll} -\operatorname{div}(|\nabla u|^{n-2}\nabla u) = \lambda \dfrac{f(u)}{|x|^\beta} & \text{in } B(1), \\ u > 0, \quad u \text{ radial in } B(1), & \\ u = 0 \quad \text{on } \partial B(1). & \end{array}\right. \tag{1.8}$$

We now have the following theorem concerning the singular Trudinger-Moser embedding:

Theorem 1.6. *Assume that* $\beta < n$ *and* f *satisfies all the conditions in Hypothesis 1, then there exists a* $\lambda_0 > 0$ *and* $\gamma_0 > 0$ *such that for all* $\lambda \in (0, \lambda_0)$, (1.8) *admits a unique solution* u *provided* $\|u\|_\infty \geq \gamma_0$.

We now give the plan of the paper: The paper is divided into four parts; In Section 2, using the Sturm's transformation, we reduce the problem to an initial value problem starting at ∞. By shooting argument, we rephrase Theorem 1.1 into the behaviour of first zero of the solution with respect to the initial condition as in [7]. We state two theorems without proof which deals with the asymptotic behaviour of the first zero and its derivative with respect to the initial data. Then we deduce the proof of Theorems 1.1 and 1.6.

In Section 3, we prove the asymptotic behaviour of first zero of the solution. Basically, this is in the ATKINSON-PELETIER [7] analysis for general γ. Here we make use of the ideas from [18] to obtain finer estimates. Since some of the estimates are used in the proof of the theorem and we were unable to find exact references, we will give the complete proof.

In Section 4, we prove the asymptotic behaviour of the derivative of the first zero with respect to the initial data using a new identity. This is much more delicate. The proof of the Theorem 1.1 follows from this finer asymptotic behaviour which will be given at the end.

2. PROOF OF THEOREM 1.1.

First we reduce the problem into an initial value problem as follows: Let $R > 0$ and define $v(x) = u\left(\dfrac{x}{R}\right)$, then, for $|x| < R$, v satisfies:

$$-\operatorname{div}(|\nabla v|^{n-2}\nabla v) = -\frac{1}{R^n}\operatorname{div}(|\nabla v|^{n-2}\nabla v)(\frac{x}{R}) = \frac{\lambda}{R^n} f(v).$$

Choosing $R = \lambda^{\frac{1}{n}}$, we see that v satisfies

$$\begin{cases} -(r^{n-1}|v'|^{n-2}v')' = f(v)r^{n-1} \quad \text{in } (0,R), \\ v > 0 \quad \text{in } (0,R), \\ v'(0) = v(R) = 0, \end{cases} \tag{2.1}$$

where v' denotes the derivatives of v with respect to $r := |x|$. Hence, for $r > 0$, consider the following initial value problem:

$$\begin{cases} -(r^{n-1}|w'|^{n-2}w')' = f(w)r^{n-1}, \\ w(0) = \gamma, w'(0) = 0. \end{cases} \tag{2.2}$$

We denote by $R(\gamma)$ to be the first zero of w defined by

$$R(\gamma) = \sup\{r \ : \ w(s) > 0 \quad \text{for all } s \in [0,r]\}.$$

Let $v(0) = \gamma$, then (2.1) and (2.2) are related by $w(0) = \gamma$, $R(\gamma) = R$, $v(r) = w(r)$ and

$$R(\gamma) = \lambda^{\frac{1}{n}}.$$

Hence the existence and uniqueness of solutions (1.7) are related to studying the behaviour of $\gamma \mapsto R(\gamma)$. Denote the solution of (2.2) by $w = w(r, \gamma)$ and now consider the Sturm's change of variables:

$$r = ne^{-\frac{t}{n}}, \quad y(t,\gamma) = w(r,\gamma), \quad R(\gamma) = ne^{-\frac{T(\gamma)}{n}},$$

then y satisfies

$$\begin{cases} -(|y'|^{n-2}y')' = f(y)e^{-t} \quad \text{in } (T(\gamma),\infty), \\ y > 0 \quad \text{in } (T(\gamma),\infty), \\ y(\infty,\gamma) = \gamma, \quad y'(\infty,\gamma) = 0, \quad y(T(\gamma),\gamma) = 0. \end{cases} \tag{2.3}$$

Before we proceed further, we prove the following important Lemma:

Lemma 2.1. *Let $f \in C^1(0,\infty) \cap C^0[0,\infty)$ such that $f \geq 0$ and $\gamma > 0$. Then, there exists a unique solution to* (2.3) *such that $\gamma \mapsto T(\gamma)$ and $\gamma \mapsto y(t,\gamma)$ are C^1 mappings. Furthermore if there exists a $p \geq 0$ and a corresponding constant $C_p > 0$ with f satisfying*

$$\overline{\lim_{\theta \to 0}} \frac{f(\theta)}{\theta^p} = C_p, \tag{2.4}$$

then there exists $\theta_0 \in \mathbb{R}$ such that

$$\begin{cases} \overline{\lim_{\gamma \to 0}}\, T(\gamma) = -\infty \quad \text{if } p > n-1, \\ \\ \overline{\lim_{\gamma \to 0}}\, T(\gamma) < \theta_0 \quad \text{if } p = n-1, \\ \\ \overline{\lim_{\gamma \to 0}}\, T(\gamma) = \infty \quad \text{if } 0 \leq p < n-1. \end{cases} \tag{2.5}$$

Proof. Consider the following initial value problem

$$\begin{cases} -(|y'|^{n-2}y')' = f(y)e^{-t} = e^{g(y)-t}, \\ y(\infty) = \gamma, \quad y'(\infty) = 0. \end{cases} \tag{2.6}$$

Local existence and uniqueness of solution to (2.6) were proved in [14]. Let $T(\gamma)$ denote the first zero of y as defined by

$$T(\gamma) := \inf\{t \ : \ y(s) > 0 \text{ for all } s \in (t,\infty)\}.$$

Integrating (2.6), we see that

$$|y'(t)|^{n-2}y'(t) = \int_t^\infty f(y(s))e^{-s}\mathrm{d}s, \tag{2.7}$$

which gives $y' > 0$ and hence we see that y must be an increasing function. We must either have $T(\gamma) > -\infty$ or $T(\gamma) = -\infty$.

Suppose $T(\gamma) = -\infty$, then $y(t) > 0$ for all $t \in \mathbb{R}$. Fixing any $t \leq 0$, we get

$$y'(t)^{n-1} = \int_t^\infty f(y(s))e^{-s}\mathrm{d}s \geq \int_0^\infty f(y(s))e^{-s}\mathrm{d}s \overset{\text{def}}{=} \beta^{n-1} > 0.$$

Integrating the above expression, we obtain $0 \leq y(t) \leq y(0)+\beta t \to -\infty$ as $t \to -\infty$ which is a contradiction. Hence $T(\gamma) \in \mathbb{R}$ and y is an increasing function in $(T(\gamma), \infty)$.

Denote the solution $y(\cdot) \overset{\text{def}}{=} y(\cdot, \gamma)$, then [14] gives $\gamma \mapsto y(t, \gamma)$ and $\gamma \mapsto T(\gamma)$ are C^1 maps.

Let f satisfy (2.4), then integrating (2.7), we obtain

$$\gamma = y(T(\gamma)) + \int_{T(\gamma)}^\infty \left(\int_\theta^\infty f(y)e^{-s}\mathrm{d}s\right)^{\frac{1}{n-1}} \mathrm{d}\theta. \tag{2.8}$$

Let $p \geq n-1$, then from (2.4), it is easy to see that there exists constants $c > 0$ and $\varepsilon < 1$ such that for all $\gamma \in (0, \varepsilon)$, the following holds:

$$f(\gamma) \leq c\gamma^p. \tag{2.9}$$

From (2.8) and (2.9), we now have

$$\gamma \leq (c\gamma^p)^{\frac{1}{n-1}} \int_{T(\gamma)}^\infty \left(\int_\theta^\infty e^{-s}\mathrm{d}s\right)^{\frac{1}{n-1}} \mathrm{d}\theta = c^{\frac{1}{n-1}}(n-1)\gamma^{\frac{p}{n-1}}e^{-\frac{T(\gamma)}{n-1}},$$

that is

$$T(\gamma) \leq \log(c(n-1)^{n-1}) + (p-(n-1))\log(\gamma) \leq \log(c(n-1)^{n-1}) \quad \text{if } p = n-1$$

$$\lim_{\gamma\to\infty} T(\gamma) \to -\infty \qquad \text{if } p > n-1.$$

Let $0 \leq p < n-1$ and suppose for some sequence $\gamma \to 0$, $T(\gamma)$ satisfies

$$\alpha := \limsup_{\gamma\to 0^+} T(\gamma) < \infty.$$

Let $m \in \mathbb{R}$ and $\beta = e^{\frac{T(\gamma)-m}{(n-1)-p}}$, then from the above assumption, β is bounded as $\gamma \to 0$. Now define w by

$$w(t) \overset{\text{def}}{=} \beta y(t - m + T(\gamma)),$$

which satisfies

$$\begin{cases} -((w')^{n-1})'(t) = \left(\dfrac{f(y(t-m+T(\gamma)))}{y^p(t-m+T(\gamma))}\right) w^{p-(n-1)}(t)w^{n-1}(t)e^{-t} & \text{in } (m,\infty) \\ w > 0 \quad \text{in } (m,\infty), \\ w(\infty) = \beta\gamma, \quad w'(\infty) = w(m) = 0. \end{cases}$$

Let φ and m_0 satisfy

$$\begin{cases} -((\varphi')^{n-1})'(t) = \varphi^{n-1}e^{-t} \quad \text{in } (m_0,\infty) \\ \varphi > 0 \quad \text{in } (m_0,\infty), \\ \varphi(\infty) = 1, \quad \varphi'(\infty) = \varphi(m_0) = 0. \end{cases}$$

Let $m = m_0 + 1$ and define $a(t) := \dfrac{f(y(t-m+T(\gamma)))}{y^p(t-m+T(\gamma))} w(t)^{p-(n-1)}$. Then from (2.4) for γ sufficiently small, we have that $a(t) \geq 1$ for $t \in (m_0, \infty)$. Since $m > m_0$ and $w > 0$ in (m, ∞), we have by the Picone's identity

$$0 \leq \int_{m_0}^{\infty} \left((\varphi')^n - \left(\frac{\varphi^n}{w^{n-1}} \right)' (w')^{n-1} \right) \mathrm{d}t = \int_{m_0}^{\infty} (1 - a(t)) \varphi^n e^{-t} \mathrm{d}t < 0$$

which is a contradiction and this proves the lemma. □

Next we consider the behaviour of $\gamma \to T(\gamma)$ as $\gamma \to \infty$ and we have the following Theorem whose proof will be given in section 3.

Theorem 2.2. *Assume that f satisfies (H1), (H2) of Hypothesis 1 and let $\beta > 0$ as in* (1.4). *Then there exists a $\gamma_0 > 0$ such that for all $\gamma > \gamma_0$, we have*

$$y'(T(\gamma), \gamma) = \frac{n}{n-1} \frac{1}{g'} + \frac{n^2 \alpha_n g''}{(n-1)(g')^3} + \mathcal{O} \left(\frac{\delta^2 g''}{(g')^4} + \frac{e^{-(g - (\frac{n-1}{n}) \gamma g')}}{g'} \right), \tag{2.10}$$

$$\begin{aligned} T(\gamma) &= (g - \left(\frac{n-1}{n} \right) \gamma g') + (n-1) \log(\left(\frac{n-1}{n} \right) g') + \frac{\alpha_n (n-1) \gamma g''}{g'} \\ &+ \mathcal{O} \left(\frac{(\log g')^2}{g'} + \frac{(\log g')^{\beta+1}}{(g')^\beta} + e^{-(g - (\frac{n-1}{n}) \gamma g')} \right) \end{aligned} \tag{2.11}$$

where $g = g(\gamma)$, $g' = g'(\gamma)$, $g'' = g''(\gamma)$, $g''' = g'''(\gamma)$, $\delta = \log(g')$ and $\alpha_n = 1 + \dfrac{1}{2} + \cdots + \dfrac{1}{n}$.

Linearization: Let $V_1(t) = V_1(t, \gamma) = \dfrac{\partial y(t, \gamma)}{\partial \gamma}$. Then differentiating (2.3) with respect to γ, we obtain:

$$\begin{cases} -((y')^{n-2} V_1')'(t) = \dfrac{f'(y(t)) V_1 e^{-t}}{n-1} & \text{in } (T(\gamma), \infty) \\ V_1(\infty) = 1, V_1'(\infty) = 0. \end{cases} \tag{2.12}$$

Differentiating $y(T(\gamma), \gamma) = 0$, we obtain

$$T'(\gamma) = -\frac{\frac{\partial y(T(\gamma), \gamma)}{\partial \gamma}}{y'(T(\gamma), \gamma)} = -\frac{V_1(T(\gamma), \gamma)}{y'(T(\gamma), \gamma)} \tag{2.13}$$

where $T'(\gamma)$ denotes the derivative of $T(\gamma)$ with respect to γ. Then using Theorem 2.2 together with the asymptotics of $V_1(T(\gamma), \gamma)$ proved in Section 4, we are able to prove the following asymptotic behaviour of $T'(\gamma)$ as $\gamma \to \infty$.

Theorem 2.3. *There exists a $\gamma_0 > 0$ such that for all $\gamma > \gamma_0$*

$$T'(\gamma) = \frac{1}{n}(g' - (n-1)\gamma g'') + \mathcal{O} \left(g'' \frac{(\log(g'))^4}{g'} \right). \tag{2.14}$$

Proof of Theorem 1.1. From (2.2), $\dfrac{\gamma g''}{g'} = \mathcal{O}(1)$ and hence from (H2) of Hypothesis 1 and (2.11), we see that $T(\gamma) \to \infty$ as $\gamma \to \infty$. That is $\lambda^{\frac{1}{n}} = R(\gamma) \to 0$ as $\gamma \to \infty$. Hence in order to prove the theorem, it is enough to show that there exists a γ_0 large such that for all $\gamma > \gamma_0$, $T(\gamma)$ is strictly increasing function. From (H3), we see that $g' - (n-1)\gamma g'' > 0$ and $\dfrac{g''}{g'}(\log(g'))^4 = o(1)(g' - (n-1)\gamma g'')$. Hence from (2.14), we have for γ_0 large and $\gamma > \gamma_0$,

$$T'(\gamma) > 0.$$

This gives that $\gamma \mapsto T(\gamma)$ is a strictly increasing function and y is nondegenerate. This proves Theorem 1.1. $\square$

Proof of Theorem 1.6. If u is a solution of (1.8), then, as earlier, converting the equation (1.8) to an initial value problem like in (2.2) we have for $R(\gamma) = \lambda^{\frac{1}{n}}$, $y(t,\gamma) = u(r,\gamma)$, $u(0) = \gamma$ and $r = ne^{-\frac{t}{n}}$; y satisfies:

$$-((y')^{n-1})' = \frac{f(y)e^{-t}}{n^\beta e^{-\frac{\beta t}{n}}} = \frac{1}{n^\beta} f(y) e^{-(1-\frac{\beta}{n})t}.$$

Let $a = (1 - \dfrac{\beta}{n})$ and $y(t) = y(\dfrac{t}{a})$, then y satisfies

$$\begin{cases} -((y')^{n-1})' = \dfrac{f(y)}{n^\beta a^n} e^{-t} \quad \text{in } (T(\gamma), \infty) \\ y(\infty) = \gamma, \ y'(\infty) = 0, \ y(T(\gamma) = 0. \end{cases}$$

Let $\tilde{f}(y) \overset{\text{def}}{=} \dfrac{f(y)}{n^\beta a^n}$, then $\tilde{g}(s) \overset{\text{def}}{=} \log(\tilde{f}(s)) = \log(f(s)) + \log(\dfrac{1}{n^\beta a^n}) = g(s) + \log(\dfrac{1}{n^\beta a^n})$. Therefore the theorem now follows from Theorem 1.1. $\square$

3. Proof of Theorem 2.2.

We shall use the following notation throughout the rest of the paper.

Notation: Let $A(\gamma)$, $B(\gamma)$ be two functions on any interval $J \subset \mathbb{R}$. We then say $A(\gamma) \sim B(\gamma)$ if there exists a constant $C > 0$ such that

$$\frac{1}{C} A(\gamma) \leq B(\gamma) \leq C A(\gamma) \quad \text{holds for all} \quad \gamma \in J. \tag{3.1}$$

Proposition 3.1. *Let $a > 0$, $q > 0$, A and B are functions on $[s_0, \infty)$ such that*

$$A(s) = as^q + B(s),$$
$$\lim_{s\to\infty} \frac{B(s)}{s^q} = 0.$$

Let $0 < \delta \leq C\log(s)$ for some $C > 0$. Then there exists an $s_1 = s_1(\delta) > 0$ and a $C_1 = C_1(\delta) > 0$ such that for all $s \geq s_1$, $\eta \in [s - \delta, s]$:

$$A(s) \sim s^q \tag{3.2}$$

$$\frac{1}{C_1} A(s) \leq A(\eta) \leq C_1 A(s).$$

Proof. (3.2) follows from (3.1). Let $\eta = s - \alpha$ with $0 \leq \alpha \leq \delta \leq C\log(s)$, and

$$A(\eta) = a(s-\alpha)^q + B(s-\alpha) = \frac{A(s)}{(1 + \frac{B(s)}{as^q})} \left(1 - \frac{\alpha}{s}\right)^q \left(1 + \frac{B(s-\alpha)}{a(s-\alpha)^q}\right).$$

Hence, $\displaystyle\lim_{s\to\infty} \frac{A(\eta)}{A(s)} = 1$ and this proves the proposition. $\square$

Proposition 3.2. *Let g satisfy (H1) and $g^{(k)}$ denote the k^{th} derivative of g. Then there exists a s_0 such that for all $\gamma > s_0$:*

(i) *$k \in \{0,1,2\}$, $g^{(k)}(\gamma) \sim \gamma^{q-k}$ and $g^{(3)}(\gamma) = \mathcal{O}(\gamma^{q-3})$,*

(ii) *Let $0 < \delta \leq C_1 \log(g'(\gamma))$, $\eta \in [\gamma - \delta, \gamma]$, $0 \leq \theta_1, \theta_2 \leq \delta$ and $k \in \{0,1,2\}$. Then there exist $C_2 = C_2(\delta)$ and $\gamma_0 = \gamma_0(\delta)$ such that for all $\gamma \geq \gamma_0$,*

$$\frac{1}{C_2} g^{(k)}(\gamma) \leq g^{(k)}(\eta) \leq C_2 g^{(k)}(\gamma),$$

and

$$\frac{\gamma g^{(k+1)}(\gamma-\theta_1)}{g^{(k)}(\gamma-\theta_2)} = \mathcal{O}(1).$$

(iii) $g^{-1}(g(\gamma)-\eta) = \gamma - \dfrac{\eta}{g'(\gamma)} + \mathcal{O}\left(\dfrac{\eta^2 g^{(2)}(\gamma)}{(g'(\gamma))^3}\right).$

Proof. (i) follows from (3.2). (ii) follows from (H1) of Hypothesis 1, (i) of Proposition 3.2 and Proposition 3.1. Differentiating $\gamma = g^{-1}(g(\gamma))$, we obtain

$$(g^{-1})'(g(\gamma)) = \frac{1}{g'(\gamma)}, \quad (g^{-1})^{(2)}(g(\gamma)) = -\frac{g^{(2)}(\gamma)}{(g'(\gamma))^3}, \quad (g^{-1})^{(2)}(\xi) = -\frac{g^{(2)}(g^{-1}(\xi))}{(g'(g^{-1}(\xi)))^3}.$$

For γ large, $g(\frac{\gamma}{2}) \le g(\gamma)-\eta$ and hence for $\xi \in [g(\gamma)-\eta, g(\gamma)]$, $\frac{\gamma}{2} \le g^{-1}(\xi) \le \gamma$. Hence, from (i) and (ii), we see that

$$(g'(g^{-1}(\xi))) \sim g'(\gamma) \quad \text{and } g^{(2)}(g^{-1}(\gamma)) \sim g^{(2)}(\gamma).$$

Therefore, by Taylor's theorem, there exists an $\xi \in [g(\gamma)-\eta, g(\gamma)]$ such that

$$\begin{aligned} g^{-1}(g(\gamma)-\eta) &= \gamma - \frac{\eta}{g'(\gamma)} + \frac{1}{2}(g^{-1})^{(2)}(\xi)\eta^2 \\ &= \gamma - \frac{\eta}{g'(\gamma)} - \frac{1}{2}\frac{g^{(2)}(g^{-1}(\xi))\eta^2}{(g'(g^{-1}(\xi)))^3} \\ &= \gamma - \frac{\eta}{g'(\gamma)} + \mathcal{O}(\frac{\eta^2 g^{(2)}(\gamma)}{(g'(\gamma))^3}). \end{aligned}$$

This proves (iii) and hence the proposition. ☐

Proposition 3.3. *Let $a > 0$ and $F(x,b) = x^n - ax^{n-1} - b$. Then there exists an $\varepsilon_0 > 0$ and a smooth function $X : (-\varepsilon_0, \varepsilon_0) \to \mathbb{R}$ such that for $|b| < \varepsilon_0$:*

$$X(0) = a,\ F(X(b), b) = 0,\ X(b) = a + \frac{b}{a^{n-1}} + \mathcal{O}\left(\frac{b^2}{a^{2n-1}}\right).$$

Proof. Since $F(a,0) = 0$ and $\dfrac{\partial F}{\partial x}(a,0) = a^{n-1} \neq 0$, hence by implicit function theorem, there exists $\varepsilon_0 > 0$ and a unique smooth function $X : (-\varepsilon_0, \varepsilon_0) \to \mathbb{R}$ such that $X(0) = a$ and $F(X(b), b) = 0$ for $|b| < \varepsilon_0$. Furthermore,

$$\begin{aligned} 0 &= \frac{\mathrm{d}}{\mathrm{d}b}F(X(b),b) = \frac{\partial F}{\partial x}(X(b),b)X'(b) + \frac{\partial F}{\partial b}(X(b),b), \\ X'(b) &= \frac{1}{nX(b)^{n-1} - (n-1)aX(b)^{n-2}} \\ X''(b) &= -\left(\frac{n(n-1)X(b)^{n-2} - (n-1)(n-2)aX(b)^{n-3}}{(nX(b)^{n-1} - (n-1)aX(b)^{n-2})^2}\right)X'(b). \end{aligned}$$

Therefore, $X'(0) = \dfrac{1}{a^{n-1}}$, $X''(0) = -\dfrac{2(n-1)}{a^{2n-1}}$ and

$$X(b) = X(0) + X'(0)b + \mathcal{O}(X''(0)b^2) = a + \frac{b}{a^{n-1}} + \mathcal{O}\left(\frac{b^2}{a^{2n-1}}\right).$$

This completes the proof of the proposition. ☐

With these preliminary propositions, we will move on to the proof of the theorem. Let $s_0 > 0$ such that g is strictly increasing convex function in $[s_0, \infty)$ and for $t \in \mathbb{R}$, $\theta \geq s_0$, $\gamma \geq s_0$ with $g^{(k)} = g^{(k)}(\gamma)$, define the following:

$$T_1 := g + (n-1)\log\left(\frac{(n-1)g'}{n}\right),$$

$$T_0 := g - \left(\frac{n-1}{n}\right)\gamma g' + (n-1)\log\left(\frac{(n-1)g'}{n}\right) - (n-1)\log\left(1 - e^{\frac{-\gamma g'}{n}}\right), \tag{3.3}$$

$$z(t) := \gamma - \frac{n}{g'}\log\left(1 + e^{\frac{T_1 - t}{n-1}}\right),$$

$$\psi(\theta) := g(\theta) - g + \left(\frac{n-1}{n}\right)(\gamma - \theta)g' - (n-1)\log\left(\frac{(n-1)g'}{n}\right). \tag{3.4}$$

Then from direct computations, z satisfies the following

$$z(T_0) = 0, \quad z(t) > 0 \ \text{ for } t > T_0, \tag{3.5}$$

$$z'(t) = \left(\frac{n}{n-1}\right)\frac{1}{g'}\frac{e^{\frac{T_1 - t}{n-1}}}{1 + e^{\frac{T_1 - t}{n-1}}} = \left(\frac{n}{n-1}\right)\frac{1}{g'}\left(1 - \frac{1}{1 + e^{\frac{T_1 - t}{n-1}}}\right), \tag{3.6}$$

$$-\left((z'(t))^{n-1}\right)' = \left(\frac{n}{(n-1)g'}\right)^{n-1}\frac{e^{T_1 - t}}{(1 + e^{\frac{T_1 - t}{n-1}})^n} = e^{g - t + g'(z - \gamma)} \tag{3.7}$$

$$z(\infty) = \gamma, \quad z'(\infty) = 0. \tag{3.8}$$

Let $\gamma > s_0$ and $y(t) = y(t, \gamma)$ be the corresponding solution of (2.6). Let $\tilde{T} > T(\gamma)$ be defined by

$$y(\tilde{T}) = s_0. \tag{3.9}$$

Then we have the following Lemma:

Lemma 3.4. *Let $\delta = \mathcal{O}(\log g')$ and*

$$T_\delta = T_1 - (n-1)\delta. \tag{3.10}$$

Then, for all $\tilde{T} \leq t$,

$$y(t) \leq z(t), \tag{3.11}$$

$$g(y(t)) \geq g - n\log\left(1 + e^{\frac{T_1 - t}{n-1}}\right) = g + g'(z - \gamma) \tag{3.12}$$

$$g(y(t)) - t \leq \psi(y(t)). \tag{3.13}$$

Furthermore, there exists a $\gamma_0 = \gamma_0(\delta) > s_0$ such that for all $\gamma \geq \gamma_0$, $t \geq T_\delta$,

$$\tilde{T} < T_\delta, \tag{3.14}$$

$$y(t) = z(t) + \mathcal{O}\left(\frac{g'' \log(1 + e^\delta)^2}{(g')^3}\right), \tag{3.15}$$

$$0 \leq g(y(t)) - g + n\log\left(1 + e^{\frac{T_1 - t}{n-1}}\right) \leq \frac{n^2 g''}{2(g')^2}\log(1 + e^{\frac{T_1 - t}{n-1}})^2 + \mathcal{O}\left(\frac{g''' \log(1 + e^\delta)^3}{(g')^3}\right) \tag{3.16}$$

$$y'(t) = z'(t)e^{\mathcal{O}\left(\frac{g'' \log(1 + e^\delta)^2}{(g')^2}\right)} \tag{3.17}$$

and

$$g'(y(t)) = g' - \frac{ng''}{g'}\left(\frac{T_1 - t}{n-1}\right) - \frac{ng''}{g'}\log\left(1 + e^{\frac{t-T_1}{n-1}}\right) + \mathcal{O}\left(\frac{g''}{(g')^2}\log(1+e^{\delta})^2\right). \tag{3.18}$$

Proof. Integrating (2.6) from t to ∞ to obtain

$$y'(t)^{n-1} = \int_t^{\infty} f(y(s))e^{-s}\mathrm{d}s. \tag{3.19}$$

Integrating (3.19) along with Fubini, we get

$$y(t) = \gamma - \int_t^{\infty} f(y(s))e^{-s}\left(\int_t^s \frac{\mathrm{d}\theta}{y'(\theta)^{n-2}}\right)\mathrm{d}s. \tag{3.20}$$

Since $\theta \mapsto g(\theta)$ and hence $\theta \mapsto f(\theta)$ are increasing functions for $\theta \geq s_0$ and hence for $t \geq \tilde{T}$, we see that

$$f(y(t))e^{-t} \leq \int_t^{\infty} f(y(s))e^{-s}\mathrm{d}s \leq f(\gamma)e^{-t}.$$

This implies

$$\lim_{t\to\infty}\left(t + (n-1)\log(y'(t))\right) = g. \tag{3.21}$$

Define

$$E(t) := y'(t)^{n-1} - \left(\frac{n-1}{n}\right)y'(t)^n g'(y(t)) - e^{g(y(t))-t}. \tag{3.22}$$

It is easy to that $E(\infty) = 0$, and now differentiating (3.22), we obtain

$$E'(t) = -\left(\frac{n-1}{n}\right)(y'(t))^{n+1}g''(y(t)) < 0 \qquad \text{for } t > \tilde{T}, \tag{3.23}$$

since $g''(\theta) > 0$ for $\theta > s_0$ by assumption. Hence, $E(t)$ is a decreasing function on $[\tilde{T}, \infty)$ and $E(t) > E(\infty) = 0$. Let $t > \tilde{T}$, then integrating (3.23), we obtain

$$0 \leq y'(t)^{n-1} - \left(\frac{n-1}{n}\right)y'(t)^n g'(y(t)) + \left(y'(t)^{n-1}\right)' = \left(\frac{n-1}{n}\right)\int_t^{\infty} y'(s)^{n+1}g''(y(s))\mathrm{d}s \tag{3.24}$$

which implies

$$\begin{aligned}
0 &\leq 1 - \left(\frac{n-1}{n}\right)y'(t)g'(y(t)) + \frac{(y'(t)^{n-1})'}{y'(t)^{n-1}} \\
&= \left(\frac{n-1}{n}\right)\int_t^{\infty}\left(\frac{y'(s)}{y'(t)}\right)^{n-1} y'(s)^2 g''(y(s))\mathrm{d}s \\
&\leq \left(\frac{n-1}{n}\right)y'(t)\int_t^{\infty} y'(s)g''(y(s))\mathrm{d}s = \left(\frac{n-1}{n}\right)y'(t)(g' - g'(y(t))).
\end{aligned}$$

We rewrite the above inequality as

$$0 \leq \left(t - \left(\frac{n-1}{n}\right)g(y(t)) + (n-1)\log y'(t)\right)' \leq \left(\frac{n-1}{n}\right)(y(t)g' - g(y(t)))',$$

and now integrating the above expression and by making use of (3.21), we obtain

$$0 \leq -\left(t - \left(\frac{n-1}{n}\right)g(y(t)) + (n-1)\log y'(t)\right) + g - \left(\frac{n-1}{n}\right)g \tag{3.25}$$

$$\leq \left(\frac{n-1}{n}\right)[(\gamma - y(t))g' - g + g(y(t)]. \tag{3.26}$$

Inequality (3.25) gives

$$y'(t)e^{-\frac{g(y(t))}{n}} \leq e^{\frac{g}{n(n-1)}-\frac{t}{n-1}}.$$

Since $g'(y(t)) \geq 0$ and increasing on $[\tilde{T}, \infty)$, we see that

$$-\left(e^{-\frac{g(y(t))}{n}}\right)' \leq \frac{g'}{n} e^{\frac{g}{n(n-1)}-\frac{t}{n-1}}.$$

Integrating the above expression from t to ∞ and simplifying, we obtain

$$g(y(t)) \geq g - n\log\left(1+\left(\frac{n-1}{n}\right)g'e^{\frac{g-t}{n-1}}\right) = g - n\log\left(1+e^{\frac{T_1-t}{n-1}}\right),$$

which proves inequality (3.12).

Now inequality (3.26) which is given by

$$y'(t)e^{\frac{g'}{n}(\gamma-y(t))} \geq e^{\frac{g-t}{n-1}}$$

after integrating from t to ∞ and simplifying, we obtain

$$y(t) \leq \gamma - \frac{n}{g'}\log\left(1+\left(\frac{n-1}{n}\right)g'e^{\frac{g-t}{n-1}}\right) = \gamma - \frac{n}{g'}\log\left(1+e^{\frac{T_1-t}{n-1}}\right)$$

which proves (3.11).

Using (3.11), it is easy to see that

$$e^{\frac{T_1-t}{n-1}} \leq e^{\frac{g'}{n}(\gamma-y(t))},$$

which after taking logarithms and simplifying, implies

$$g(y(t)) - t \leq \psi(y(t))$$

and this completes the proof of (3.13).

Proof of inequality (3.14) will be by contradiction, suppose $T_\delta \leq \tilde{T}$, then $\dfrac{T_1-\tilde{T}}{n-1} \leq \delta$, then (3.12) gives

$$g(s_0) = g(y(\tilde{T})) \geq g - n\log\left(1+e^{\frac{T_1-\tilde{T}}{n-1}}\right) \geq g + \mathcal{O}(\log(g')) \to \infty \quad \text{as } \gamma \to \infty,$$

which is a contradiction and hence there exists a $\gamma_0 = \gamma_0(\delta) > 0$ such that inequality (3.14) holds for all $\gamma > \gamma_0$.

Let $t > T_\delta$, then $\dfrac{T_1-t}{n-1} \leq \delta$ which also implies that $\log\left(1+e^{\frac{T_1-t}{n-1}}\right) \leq \log\left(1+e^{\delta}\right)$. Let $\gamma \geq \gamma_0(\delta)$, then from (3.12), (3.13), (3.14) and setting $\eta = n\log\left(1+e^{\frac{T_1-t}{n-1}}\right)$ in assertion (iii) of Proposition 3.2, we have

$$\begin{aligned} y(t) &\geq g^{-1}\left(g - n\log(\left(1+e^{\frac{T_1-t}{n-1}}\right))\right) \\ &= \gamma - \frac{n}{g'}\log\left(1+e^{\frac{T_1-t}{n-1}}\right) + \mathcal{O}\left(\frac{g''}{(g')^3}\log(1+e^{\delta})^2\right) \\ &= z(t) + \mathcal{O}\left(\frac{g''}{(g')^3}\log(1+e^{\delta})^2\right). \end{aligned}$$

This inequality along with (3.11) proves (3.15).

From Taylor's series, (3.11) and from Proposition 3.2, there exists $\xi \in \left[\gamma - \frac{n}{g'} \log\left(1 + e^{\frac{T_1 - t}{n-1}}\right), \gamma\right]$ such that

$$\begin{aligned} g(y(t)) &\leq g(z(t)) \\ &= g - n\log\left(1 + e^{\frac{T_1-t}{n-1}}\right) + \frac{g'' n^2 \log\left(1 + e^{\frac{T_1-t}{n-1}}\right)^2}{2(g')^2} + \frac{g'''(\xi) n^3 \log\left(1 + e^{\frac{T_1-t}{n-1}}\right)^3}{3!(g')^3} \\ &= g - n\log\left(1 + e^{\frac{T_1-t}{n-1}}\right) + \frac{g'' n^2 \log\left(1 + e^{\frac{T_1-t}{n-1}}\right)^2}{2(g')^2} + \mathcal{O}\left(\frac{g'''}{(g')^3} \log\left(1 + e^{\delta}\right)^3\right). \end{aligned}$$

The above inequality together with (3.12) proves (3.16).

From (3.12), it is easy to see that

$$y'(t)^{n-1} = \int_t^\infty e^{g(y(s)) - s} \mathrm{d}s \geq \int_t^\infty e^{g + g'(z(s) - \gamma) - s} \mathrm{d}s = z'(t)^{n-1}. \tag{3.27}$$

Inequality (3.16) implies

$$\begin{aligned} y'(t)^{n-1} &= \int_t^\infty e^{g(y(s)) - s} \mathrm{d}s \\ &\leq e^{\mathcal{O}\left(\frac{g''}{(g')^2} \log\left(1+e^\delta\right)^2\right)} \int_t^\infty e^{g + g'(z - \gamma) - s} \mathrm{d}s \\ &= e^{\mathcal{O}\left(\frac{g''}{(g')^2} \log\left(1+e^\delta\right)^2\right)} z'(t)^{n-1} \end{aligned}$$

and this proves (3.17).

From (3.15), we have

$$\begin{aligned} g'(y(t)) &= g'\left(\gamma - \frac{n}{g'}\log\left(1 + e^{\frac{T_1-t}{n-1}}\right) + \mathcal{O}\left(\frac{g''}{(g')^3}\log\left(1+e^\delta\right)^2\right)\right) \\ &= g' - \frac{ng''}{g'}\log\left(1 + e^{\frac{T_1-t}{n-1}}\right) + \mathcal{O}\left(\frac{g''}{(g')^2}\log\left(1+e^\delta\right)^2\right) \\ &= g' - \frac{ng''}{g'}\left(\frac{T_1 - t}{n-1}\right) - \frac{ng''}{g'}\log\left(1 + e^{\frac{t-T_1}{n-1}}\right) + \mathcal{O}\left(\frac{g''}{(g')^2}\log\left(1+e^\delta\right)^2\right) \end{aligned}$$

which proves (3.18) and this completes the proof of the lemma. ☐

Lemma 3.5. *Let $k \geq 1$ be an integer and $X(t) \overset{\text{def}}{=} e^{\frac{T_1-t}{n-1}}$. Then the following statements hold:*

$$\alpha_k := 1 + \frac{1}{2} + \frac{1}{3} + \ldots + \frac{1}{k} = -\sum_{r=1}^{k} \frac{(-1)^r C_r^k}{r} \tag{3.28}$$

$$\sum_{r=0}^{k} \frac{(-1)^r C_r^k}{(r+2)} = \frac{1}{(k+1)(k+2)}, \tag{3.29}$$

$$\int_t^\infty (z'(s))^{k+1} \mathrm{d}s = \left(\frac{n}{(n-1)g'}\right)^{k+1} \left[-\alpha_k + \log(1 + X(t)) - \sum_{r=1}^{k} \frac{(-1)^r C_r^k}{r(1 + X(t))^r}\right]. \tag{3.30}$$

Proof. By explicit integration, we get

$$\alpha_k = \sum_{r=0}^{k-1} \int_0^1 x^r \mathrm{d}x = \int_0^1 \frac{1 - x^k}{1 - x} \mathrm{d}x = \int_0^1 \frac{1 - (1-x)^k}{x} \mathrm{d}x = -\sum_{r=1}^{k} \frac{(-1)^r C_r^k}{r}.$$

Similar calculation gives

$$\sum_{r=0}^{k}\frac{(-1)^r C_r^k}{(r+2)}=\int_0^1(1-x)^k x\mathrm{d}x=\int_0^1 x^k(1-x)\mathrm{d}x=\frac{1}{(k+1)(k+2)}.$$

Since $\mathrm{d}X(s)=-\dfrac{X(s)\mathrm{d}s}{(n-1)}$, making use of (3.28), we get

$$\begin{aligned}\int_t^\infty z'(s)^{k+1}\mathrm{d}s &= \left(\frac{n}{(n-1)g'}\right)^{k+1}\int_t^\infty \frac{X(s)^{k+1}}{(1+X(s))^{k+1}}\mathrm{d}s\\ &= \left(\frac{n}{(n-1)g'}\right)^{k+1}(n-1)\int_0^{X(t)}\frac{x^k}{(1+x)^{k+1}}\mathrm{d}x\\ &= \left(\frac{n}{(n-1)g'}\right)^{k+1}(n-1)\sum_{r=0}^{k}(-1)^r C_r^k\int_0^{X(t)}\frac{\mathrm{d}x}{(1+x)^{r+1}}\\ &= \left(\frac{n}{(n-1)g'}\right)^{k+1}(n-1)\left[\log(1+x)-\sum_{r=1}^{k}\frac{(-1)^r C_r^k}{r(1+x)^r}\right]_0^{X(t)}\\ &= \left(\frac{n}{(n-1)g'}\right)^{k+1}(n-1)\left[-\alpha_k+\log(1+X(t))-\sum_{r=1}^{k}\frac{(-1)^r C_r^k}{r(1+X(t))^r}\right].\end{aligned}$$

This completes the proof of the lemma. ☐

Lemma 3.6. *Let $k\geq 3$, $\delta=k\log(g')$ and $T_\delta=T_1-(n-1)\delta$. Then there exists a $\gamma_0=\gamma_0(k)$ such that for all $\gamma\geq\gamma_0$,*

$$y'(T_\delta)=\frac{n}{(n-1)g'}+\frac{n^2\alpha_n g''}{(n-1)(g')^3}+\mathcal{O}\left(\frac{g''\delta^2}{(g')^4}+\frac{1}{(g')^{k+1}}\right) \tag{3.31}$$

with α_n defined as in (3.28).

Proof. To simplify the notation for the proof of this lemma, denote $y=y(T_\delta)$, $y'=y'(T_\delta)$, $z=z(T_\delta)$ and $z'=z'(T_\delta)$, then from (3.24) we have

$$\begin{aligned}(y')^{n-1}-\left(\frac{n-1}{n}\right)(y')^n g' &= e^{g(y)-T_\delta}-\left(\frac{n-1}{n}\right)(g'-g'(y))(y')^n\\ &\quad+\left(\frac{n-1}{n}\right)\int_{T_\delta}^\infty g''(y(s))(y'(s)^{n+1}-(z'(s))^{n+1})\mathrm{d}s\\ &\quad+\left(\frac{n-1}{n}\right)\int_{T_\delta}^\infty z'(s)^{n+1}(g''(y(s))-g'')\mathrm{d}s\\ &\quad+\left(\frac{n-1}{n}\right)g''\int_{T_\delta}^\infty z'(s)^{n+1}\mathrm{d}s\overset{\mathrm{def}}{=}J_1+J_2+J_3+J_4+J_5.\end{aligned}$$

Since $\log\left(1+e^{\frac{T_1-T_\delta}{n-1}}\right)=\log(1+e^\delta)=\delta+\log(1+e^{-\delta})=\delta+\mathcal{O}\left(\dfrac{1}{(g')^k}\right)$ and $X(T_\delta)=e^\delta=(g')^k$, we have from (3.7) and (3.16)

$$\begin{aligned}J_1=e^{g(y)-T_\delta} &= e^{\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right)}e^{g+g'(z-\gamma)-T_\delta}\\ &= \mathcal{O}\left(((z')^{n-1})'\right)=\mathcal{O}\left(\frac{1}{(g')^{n-1}}\frac{X(T_\delta)^{n-1}}{(1+X(T_\delta))^n}\right)=\mathcal{O}\left(\frac{1}{(g')^{k+n-1}}\right).\end{aligned}$$

From (3.17) and (3.18) we have

$$\begin{aligned}
y'(T_\delta)^n &= z'(T_\delta)^n e^{\mathcal{O}\left(\frac{g''}{(g')^2}\delta^2\right)} \\
&= \left(\frac{n}{(n-1)g'}\right)^n \left(1-\frac{1}{1+X(T_\delta)}\right)^n e^{\mathcal{O}\left(\frac{g''}{(g')^2}\delta^2\right)} \\
&= \left(\frac{n}{(n-1)g'}\right)^n \left(1+\mathcal{O}(\frac{1}{(g')^k})\right) e^{\mathcal{O}\left(\frac{g''}{(g')^2}\delta^2\right)} \\
&= \left(\frac{n}{(n-1)g'}\right)^n \left(1+\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}+\frac{1}{(g')^k}\right)\right)
\end{aligned}$$

and

$$\begin{aligned}
J_2 &= \left(\frac{n-1}{n}\right)(y')^n(g'(y)-g') \\
&= \left(\frac{n-1}{n}\right)\left(\frac{n}{(n-1)g'}\right)^n (1+\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right))\left(-\frac{ng''}{g'}\delta+\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right)\right) \\
&= -\left(\frac{n}{n-1}\right)^{n-1}\frac{ng''\delta}{(g')^{n+1}}+\mathcal{O}\left(\frac{g''\delta^2}{(g')^{n+2}}\right).
\end{aligned}$$

From (3.17) and (3.30),

$$\begin{aligned}
J_3 &= \left(\frac{n-1}{n}\right)\int_{T_\delta}^{\infty}(y'(s)^{n+1}-z'(s)^{n+1})g''(y(s))\mathrm{d}s \\
&= \mathcal{O}\left(\frac{(g'')^2\delta^2}{(g')^2}\int_{T_\delta}^{\infty} z'(s)^{n+1}\mathrm{d}s\right)=\mathcal{O}\left(\frac{(g'')^2\delta^3}{(g')^{n+3}}\right).
\end{aligned}$$

From Proposition 3.2 and (3.15), for $s \geq T_\delta$, there exist a $\xi \in [y(s),\gamma]$ such that

$$\begin{aligned}
g''(y(s)) &= g''+g'''(\xi)(y(s)-\gamma) = g''+\mathcal{O}(g'''(z(s)-\gamma))+\mathcal{O}\left(\frac{g'''g''\delta^2}{(g')^3}\right) \\
&= g''+\mathcal{O}\left(\frac{g'''\delta}{g'}\right) = g''+\mathcal{O}\left(\frac{\gamma g'''}{g''}\frac{g''\delta}{\gamma g'}\right) = g''+\mathcal{O}\left(\frac{g''\delta}{(g')^{1+\frac{1}{q-1}}}\right).
\end{aligned}$$

Hence from (3.30)

$$J_4 = \left(\frac{n-1}{n}\right)\int_{T_\delta}^{\infty} z'(s)^{n+1}(g''(y(s))-g'')\mathrm{d}s = \mathcal{O}\left(\frac{g''\delta^2}{(g')^{n+2}}\right).$$

From (3.30)

$$\begin{aligned}
J_5 &= \left(\frac{n-1}{n}\right)g''\int_{T_\delta}^{\infty} z'(s)^{n+1}\mathrm{d}s \\
&= (n-1)\left(\frac{n}{n-1}\right)^n\frac{g''}{(g')^{n+1}}\left(-\alpha_n+\delta+\mathcal{O}\left(\frac{1}{(g')^k}\right)\right) \\
&= n\left(\frac{n}{n-1}\right)^{n-1}\frac{g''}{(g')^{n+1}}(-\alpha_n+\delta+\mathcal{O}\left(\frac{1}{(g')^k}\right)).
\end{aligned}$$

Combining all the estimates to obtain

$$\begin{aligned}(y')^{n-1}-\left(\frac{n-1}{n}\right)g'(y')^{n} &= J_1+J_2+J_3+J_4+J_5\\ &= -\left(\frac{n}{n-1}\right)^{n-1}\frac{ng''\delta}{(g')^{n+1}}\\ &\quad+\left(\frac{n}{n-1}\right)^{n-1}\frac{ng''}{(g')^{n+1}}\left(-\alpha_n+\delta+\mathcal{O}\left(\frac{1}{(g')^{k}}\right)\right)\\ &\quad+\mathcal{O}\left(\frac{g''\delta^2}{(g')^{n+2}}+\frac{1}{(g')^{k+n-1}}\right)\end{aligned}$$

Therefore,

$$(y')^{n}-\frac{n}{(n-1)g'}(y')^{n-1}-n\left(\frac{n}{n-1}\right)^{n}\frac{\alpha_n g''}{(g')^{n+2}}=\mathcal{O}\left(\frac{g''\delta^2}{(g')^{n+3}}+\frac{1}{(g')^{k+n}}\right)$$

and Proposition 3.3 now gives

$$y'=\frac{n}{(n-1)g'}+\frac{n^2\alpha_n g''}{(n-1)(g')^3}+\mathcal{O}\left(\frac{g''\delta^2}{(g')^4}+\frac{1}{(g')^{k+1}}\right).$$

This completes the proof of the lemma. ☐

Lemma 3.7. *Let* $k\geq 3$, $\delta=k\log(g')$, $T_\delta=T_1-(n-1)\delta$ *and* $b\in\mathbb{R}$. *Assume that* g *satisfies*

$$\limsup_{s\to\infty}(g(s)-\left(\frac{n-1}{n}\right)sg'(s))\geq b+1. \tag{3.32}$$

Then, there exists a $\gamma_0(k)>0$ *such that for all* $\gamma\geq\gamma_0(k)$, *we have*

$$\int_{\tilde{T}}^{T_\delta}e^{g(y(s))-s}\mathrm{d}s=\mathcal{O}\left(\frac{(g')^{\frac{q}{q-1}}}{(g')^{k+n-1}}\right), \tag{3.33}$$

$$\int_{\tilde{T}}^{T_\delta}g'(y(s))e^{g(y(s))-s}\mathrm{d}s=\mathcal{O}\left(\frac{(g')^{\frac{q}{q-1}}}{(g')^{k+n-2}}\right)$$

where q *is given as in* (1.3).

Proof. Choose γ_1 such that Lemma 3.4 holds for $\gamma\geq\gamma_1$, $\delta=k\log(g')$ and

$$g-\left(\frac{n-1}{n}\right)\gamma g'\geq b. \tag{3.34}$$

Let ψ be as in (3.4), then, for $\theta\geq s_0$, we have $\theta\to\psi(\theta)$ is a convex function. Hence from (3.13), we have

$$g(y(t))-t\leq\psi(y(t))\leq\max\left\{\psi(y(\tilde{T})),\psi(y(T_\delta))\right\}\qquad\text{for } t\in[\tilde{T},T_\delta]. \tag{3.35}$$

Estimate for $\psi(y(\tilde{T}))$*:* From (3.34),

$$\begin{aligned}\psi(y(\tilde{T})) &= g(s_0)-\left(\frac{n-1}{n}\right)s_0g'-(g-\left(\frac{n-1}{n}\right)\gamma g')-(n-1)\log(\left(\frac{n-1}{n}\right)g')\\ &\leq g(s_0)-\left(\frac{n-1}{n}\right)s_0g'-b-(n-1)\log(\left(\frac{n-1}{n}\right)g'),\end{aligned}$$

which implies that

$$e^{\psi(y(\tilde{T}))}=\mathcal{O}\left(\frac{e^{-\left(\frac{n-1}{n}\right)s_0g'}}{(g')^{n-1}}\right). \tag{3.36}$$

Estimate for $\psi(y(T_\delta))$: From (3.15) and (3.16)

$$\begin{aligned}
\psi(y(T_\delta)) &= g(y(T_\delta)) - g + \left(\frac{n-1}{n}\right)(\gamma - y(T_\delta))g' - (n-1)\log(\left(\frac{n-1}{n}\right)g') \\
&= -n\log(1+e^\delta) + \left(\frac{n-1}{n}\right)(\gamma - z(T_\delta))g' - (n-1)\log(\left(\frac{n-1}{n}\right)g') + \mathcal{O}\left(\frac{\delta^2}{g'}\right) \\
&= -n\log(1+e^\delta) + (n-1)\log(1+e^\delta) - (n-1)\log(\left(\frac{n-1}{n}\right)g') + \mathcal{O}\left(\frac{\delta^2}{g'}\right) \\
&= -\delta - (n-1)\log(g') + \mathcal{O}(1).
\end{aligned}$$

Thus,

$$e^{\psi(y(T_\delta))} = \mathcal{O}\left(e^{-\delta-(n-1)\log(g')}\right) = \mathcal{O}\left(\frac{1}{(g')^{k+n-1}}\right). \tag{3.37}$$

Estimate for $\tilde{T} - T_\delta$: From (3.5) and (3.11), it is easy to see that $T_0 < \tilde{T}$. Now from (3.3) and (3.14), we have for for $\gamma \geq \gamma_0$,

$$\begin{aligned}
0 < T_\delta - \tilde{T} \leq T_\delta - T_0 &= g + (n-1)\log(\left(\frac{n-1}{n}\right)g') - (n-1)\delta \\
&\quad -g - (n-1)\log(\left(\frac{n-1}{n}\right)g') + \left(\frac{n-1}{n}\right)\gamma g' + \mathcal{O}\left(e^{-\frac{\gamma g'}{n}}\right) \\
&\leq \mathcal{O}\left((g')^{\frac{q}{q-1}}\right),
\end{aligned} \tag{3.38}$$

where we have used $\gamma = \mathcal{O}\left((g')^{\frac{1}{q-1}}\right)$.

Hence from (3.36), (3.37) and (3.38), we have for γ large

$$\begin{aligned}
\int_{\tilde{T}}^{T_\delta} e^{g(y(s))-s}\mathrm{d}s &\leq (T_\delta - \tilde{T})\max\left\{e^{\psi(y(\tilde{T}))}, e^{\psi(y(T_\delta))}\right\} \\
&= \mathcal{O}\left(\frac{(g')^{\frac{q}{q-1}}}{(g')^{k+n-1}}\right)
\end{aligned}$$

and

$$\int_{\tilde{T}}^{T_\delta} g'(y(s))e^{g(y(s))-s}\mathrm{d}s = \mathcal{O}\left(g'\int_{\tilde{T}}^{T_\delta} e^{g(y(s))-s}\mathrm{d}s\right) = \mathcal{O}\left(\frac{(g')^{\frac{q}{q-1}}}{(g')^{k+n-2}}\right).$$

This completes the proof of the lemma. □

Immediate consequence of the above lemma gives the following asymptotics:

Lemma 3.8. *Let g satisfies (3.32) with $k \geq 5 + \dfrac{q}{q-1}$, $\delta = k\log(g')$ and $T_\delta = T_1 - (n-1)\delta$. Then there exists a $\gamma_0 > 0$ such that for $\gamma > \gamma_0$ and for all $t \in [\tilde{T}, T_\delta]$, we have*

$$y'(t) = \frac{n}{(n-1)g'} + \frac{n^2\alpha_n g''}{(n-1)(g')^3} + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^4}\right), \tag{3.39}$$

$$\begin{aligned}
\tilde{T} &= \left(\frac{n-1}{n}\right)s_0 g' + (g - \left(\frac{n-1}{n}\right)\gamma g') + (n-1)\log(\left(\frac{n-1}{n}\right)g') + \\
&\quad + \frac{(n-1)g''\alpha_n(\gamma - s_0)}{g'} + \mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right),
\end{aligned} \tag{3.40}$$

$$\begin{aligned} T(\gamma) &\geq \left(g - \left(\frac{n-1}{n}\right)\gamma g'\right) + (n-1)\log\left(\left(\frac{n-1}{n}\right)g'\right) + \frac{(n-1)\alpha_n\gamma g''}{g'} + \\ &+\mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right). \end{aligned} \tag{3.41}$$

Furthermore, setting $t_0 \overset{\text{def}}{=} (n+3)\log(g')$, *if* $T(\gamma) \geq t_0 + \mathcal{O}(1)$ *then*

$$\begin{aligned} T(\gamma) = g - \left(\frac{n-1}{n}\right)\gamma g' + (n-1)\log(\left(\frac{n-1}{n}\right)g') &+ \frac{(n-1)\alpha_n\gamma g''}{g'} + \\ &+ \mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right). \end{aligned} \tag{3.42}$$

Also for all $t \in [\max\{t_0, T(\gamma)\}, \tilde{T}]$, *we have*

$$y'(t) = y'(\tilde{T}) + \mathcal{O}\left(\frac{g''\delta^2}{(g')^4}\right). \tag{3.43}$$

If $T(\gamma) < t_0 + \mathcal{O}(1)$, *then*

$$y(t_0) = \mathcal{O}\left(\frac{\log(g')}{g'}\right). \tag{3.44}$$

Proof. Since $k \geq 5 + \dfrac{q}{q-1}$, we have for $t \in [\tilde{T}, T_\delta]$ and from using (3.33), (3.31)

$$\begin{aligned} y'(t)^{n-1} &= y'(T_\delta)^{n-1} + \int_t^{T_\delta} e^{g(y(s))-s}\mathrm{d}s \\ &= y'(T_\delta)^{n-1} + \mathcal{O}\left(\frac{1}{(g')^{4+n}}\right) \\ &= y'(T_\delta)^{n-1}\left(1 + \mathcal{O}\left(\frac{1}{(g')^5}\right)\right). \end{aligned}$$

Now by making use of estimate (3.31), we get

$$\begin{aligned} y'(t) &= y'(T_\delta)\left(1 + \mathcal{O}\left(\frac{1}{(g')^5}\right)\right) \\ &= \frac{n}{(n-1)g'} + \frac{n^2\alpha_n g''}{(n-1)(g')^3} + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^4}\right). \end{aligned}$$

This proves (3.39).

From the mean value theorem and (3.39), there exists a $\xi \in [\tilde{T}, T_\delta]$ such that

$$\begin{aligned}
\tilde{T} &= T_\delta + \frac{y(\tilde{T}) - y(T_\delta)}{y'(\xi)} = T_\delta + \frac{s_0 - z(T_\delta) + \mathcal{O}\left(\frac{g''\delta^2}{(g')^3}\right)}{y'(\xi)} \\
&= T_\delta + \left(\frac{n-1}{n}\right) g' \left[1 - \frac{n\alpha_n g''}{(g')^2} + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^3}\right)\right] \times \left[s_0 - \gamma + \frac{n}{g'}\log(1+e^\delta) + \mathcal{O}\left(\frac{g''\delta^2}{(g')^3}\right)\right] \\
&= T_1 - (n-1)\delta + \left(\frac{n-1}{n}\right) g'(s_0 - \gamma) + (n-1)\delta + (n-1)\log(1+e^{-\delta}) \\
&\quad -\frac{\alpha_n(n-1)g''(s_0-\gamma)}{g'} - \frac{n(n-1)\alpha_n g''}{(g')^2}\log(1+e^\delta) + \mathcal{O}\left(\frac{\delta^2\gamma g''}{(g')^2}\right) \\
&= \left(\frac{n-1}{n}\right) s_0 g' + (g - \left(\frac{n-1}{n}\right)\gamma g') + (n-1)\log(\left(\frac{n-1}{n}\right) g') + \frac{\alpha_n(n-1)g''(\gamma - s_0)}{g'} \\
&\quad +\mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right).
\end{aligned}$$

This proves (3.40).

For $t \in [T(\gamma), \tilde{T}]$, we have $y'(t) \geq y'(\tilde{T})$ and hence from the mean value theorem, there exist a $\xi \in [T(\gamma), \tilde{T}]$ such that

$$\tilde{T} - T(\gamma) = \frac{y(\tilde{T}) - y(T(\gamma))}{y'(\xi)} \leq \frac{s_0}{y'(\tilde{T})}.$$

Hence from (3.39) and (3.40)

$$\begin{aligned}
T(\gamma) &\geq \tilde{T} - \frac{s_0}{y'(\tilde{T})} \\
&= \left(\frac{n-1}{n}\right) s_0 g' + (g - \left(\frac{n-1}{n}\right)\gamma g') + (n-1)\log(\left(\frac{n-1}{n}\right) g') + \mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right) \\
&\quad + \frac{\alpha_n(n-1)g''(\gamma - s_0)}{g'} - s_0\left(\frac{n-1}{n}\right) g' \left[1 - \frac{n\alpha_n g''}{(g')^2} + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^3}\right)\right] \\
&= (g - \left(\frac{n-1}{n}\right)\gamma g') + (n-1)\log(\left(\frac{n-1}{n}\right) g') + \frac{\alpha_n(n-1)\gamma g''}{g'} \\
&\quad +\mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right). \qquad (3.45)
\end{aligned}$$

This proves (3.41).

Let $t \in [\max\{t_0, T(\gamma)\}, \tilde{T}]$, then

$$\begin{aligned}
y'(t)^{n-1} &= y'(\tilde{T})^{n-1} + \int_t^{\tilde{T}} f(y(s))e^{-s}\mathrm{d}s \\
&= y'(\tilde{T})^{n-1} + \mathcal{O}(e^{-T(\gamma)}) = y'(\tilde{T})^{n-1} + \mathcal{O}\left(\frac{1}{(g')^{n+3}}\right).
\end{aligned}$$

Hence

$$y'(t) = y'(\tilde{T}) + \mathcal{O}\left(\frac{g''\delta^2}{(g')^4}\right). \qquad (3.46)$$

Suppose that $T(\gamma) \geq t_0 + \mathcal{O}(1)$, then there exists $\xi \in [T(\gamma), \tilde{T}]$ and from (3.45) and (3.46) we have

$$\begin{aligned} T(\gamma) &= \tilde{T} + \frac{y(T(\gamma)) - y(\tilde{T})}{y'(\xi)} \\ &= \tilde{T} - \frac{s_0}{y'(\tilde{T})}(1 + \mathcal{O}\left(\frac{g''\delta^2}{(g')^3}\right)) \\ &= (g - \left(\frac{n-1}{n}\right)\gamma g') + (n-1)\log(\left(\frac{n-1}{n}\right)g') + \frac{\alpha_n(n-1)\gamma g''}{g'} + \mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right). \end{aligned}$$

This proves (3.42).

Let $T(\gamma) < t_0 = (n+3)\log(g') + \mathcal{O}(1)$. Then (3.46) holds for $t \in [t_0, \tilde{T}]$ and this proves (3.43). From (3.43) and mean value theorem, there exists a $\xi \in [t_0, \tilde{T}] \subset [T(\gamma), \tilde{T}]$ such that

$$\begin{aligned} y(t_0) &= y(\tilde{T}) + y'(\xi)(t_0 - \tilde{T}) \\ &= s_0 + y'(\tilde{T})(t_0 - \tilde{T}) + O\left(\frac{(t_0 - \tilde{T})g''\delta^2}{(g')^4}\right) \\ &= s_0 + (t_0 - \tilde{T})\left(\frac{n}{(n-1)g'} + \frac{\alpha_n n^2 g''}{(n-1)(g')^3} + \mathcal{O}\left(\frac{g''\delta^2}{(g')^4}\right)\right) \\ &= s_0 + \frac{n}{(n-1)g'}\left(1 + \mathcal{O}\left(\frac{g''}{(g')^2}\right)\right)\left(t_0 - \left(\frac{n-1}{n}\right)s_0 g'\right. \\ &\quad \left. -(g - \left(\frac{n-1}{n}\right)\gamma g') - (n-1)\log(\left(\frac{n-1}{n}\right)g') + \mathcal{O}(1)\right) \\ &= \frac{n}{(n-1)g'}\left[t_0 - (g - \left(\frac{n-1}{n}\right)\gamma g') - (n-1)\log(\left(\frac{n-1}{n}\right)g')\right] + \mathcal{O}\left(\frac{1}{g'}\right) \\ &\leq \frac{n}{(n-1)g'}\left[t_0 - b - (n-1)\log(\left(\frac{n-1}{n}\right)g')\right] + \mathcal{O}\left(\frac{1}{g'}\right) = \mathcal{O}\left(\frac{\log(g')}{g'}\right). \end{aligned}$$

This proves (3.44) and hence the lemma. ☐

Lemma 3.9. *Assume that there exists a* $\beta > 0$ *such that as* $s \to 0$,

$$f(s) - f(0) = \mathcal{O}(s^\beta).$$

Then there exists a bounded function $A : \gamma \to A(\gamma)$ *such that* $A(\gamma) = 0$ *if* $f(0) = 0$ *and for large* γ

$$\begin{aligned} T(\gamma) &= (g - \left(\frac{n-1}{n}\right)\gamma g') + (n-1)\log(\left(\frac{n-1}{n}\right)g') + \frac{(n-1)\alpha_n\gamma g''}{g'} + A(\gamma) \\ &\quad + \mathcal{O}\left(\frac{(\log(g'))^2}{g'} + \frac{(\log(g'))^{\beta+1}}{(g')^\beta}e^{-(g-(\frac{n-1}{n})\gamma g')}\right). \end{aligned} \tag{3.47}$$

Furthermore,

(i) *Assume that* $\lim_{\gamma\to\infty}(g - \left(\frac{n-1}{n}\right)\gamma g') = \infty$, *then*

$$y'(T(\gamma)) = \frac{n}{(n-1)g'} + \frac{n^2\alpha_n g''}{(n-1)(g')^3} + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^4} + \frac{e^{-(g-(\frac{n-1}{n})\gamma g')}}{(g')^{n-1}}\right). \tag{3.48}$$

(ii) *Assume that* $\limsup\limits_{\gamma\to\infty}\left(g-\left(\frac{n-1}{n}\right)\gamma g'\right)<\infty$. *Then,*

$$\begin{aligned}
y'(T(\gamma)) &= \frac{n}{(n-1)g'}+\frac{n^2\alpha_n g''}{(n-1)(g')^3}+\left[\left(1+\frac{f(0)e^{-T(\gamma)}}{y'(t_0)^{n-1}}\right)^{\frac{1}{n-1}}-1\right]y'(\tilde{T})\\
&\quad +\mathcal{O}\left(\frac{(\log(g'))^\beta}{(g')^\beta}+\frac{g''(\log(g'))^2}{(g')^4}\right). \qquad (3.49)
\end{aligned}$$

Proof. Let $t_0=(n+3)\log(g')$. If $T(\gamma)\geq t_0$, then (3.47) and (3.48) follows from Lemma 3.8.

Hence assume that $T(\gamma)<t_0$. From (3.41) it is easy to see that $(g-\frac{(n-1)}{n}\gamma g')=\mathcal{O}(\log(g'))$ which gives $\tilde{T}=\mathcal{O}(g')$ and from (3.46), we have

(3.50)

$$\begin{aligned}
t_0-\frac{y(t_0)}{y'(t_0)} &= t_0-\frac{y(\tilde{T})+y'(\xi)(t_0-\tilde{T})}{y'(\tilde{T})+\mathcal{O}\left(\frac{g''\delta^2}{(g')^4}\right)}\\
&= t_0-\frac{s_0+y'(\tilde{T})(t_0-\tilde{T})+\mathcal{O}\left(\frac{(t_0-\tilde{T})g''\delta^2}{(g')^4}\right)}{y'(\tilde{T})(1+\mathcal{O}(\frac{g''\delta^2}{(g')^3}))}\\
&= t_0-\frac{s_0}{y'(\tilde{T})}-(t_0-\tilde{T})+\mathcal{O}\left(\frac{g''\delta^2}{(g')^3}\right)\\
&= \tilde{T}-\frac{s_0}{y'(\tilde{T})}+\mathcal{O}\left(\frac{g''\delta^2}{(g')^3}\right)\\
&= (g-\left(\frac{n-1}{n}\right)\gamma g')+(n-1)\log(\left(\frac{n-1}{n}\right)g')+\frac{(n-1)\alpha\gamma g''}{g'}+\mathcal{O}\left(\frac{(\log(g'))^2}{g'}\right).
\end{aligned}$$

From (3.41), (3.39) and (3.46) we have for $t\in[T(\gamma),t_0]$,

$$\frac{e^{-t}}{y'(t_0)^{n-1}}=\mathcal{O}\left(e^{-(g-\frac{n-1}{n}\gamma g')}\right). \qquad (3.51)$$

Hence from (3.44) and (3.51), we have

$$\begin{aligned}
y'(t)^{n-1} &= y'(t_0)^{n-1}+\int_t^{t_0}f(y(s))e^{-s}\mathrm{d}s\\
&= y'(t_0)^{n-1}+f(0)e^{-t}+\mathcal{O}\left(\int_t^{t_0}y(s)^\beta e^{-s}\mathrm{d}s+\frac{1}{(g')^{n+3}}\right)\\
&= y'(t_0)^{n-1}+f(0)e^{-t}+\mathcal{O}\left(\frac{(\log(g'))^\beta}{(g')^\beta}e^{-t}+\frac{1}{(g')^{n+3}}\right).
\end{aligned}$$

Therefore, from (3.51)

$$\begin{aligned}
\frac{y'(t)}{y'(t_0)} &= \left(1+\frac{f(0)e^{-t}}{y'(t_0)^{n-1}}\right)^{\frac{1}{n-1}}+\mathcal{O}\left(\frac{(\log(g'))^\beta}{(g')^\beta}\frac{e^{-t}}{y'(t_0)^{n-1}}+\frac{1}{(g')^3}\right)\\
&= \left(1+\frac{f(0)e^{-t}}{y'(t_0)^{n-1}}\right)^{\frac{1}{n-1}}+\mathcal{O}\left(\frac{(\log(g'))^\beta}{(g')^\beta}e^{-(g-\frac{n-1}{n}\gamma g')}+\frac{1}{(g')^3}\right). \qquad (3.52)
\end{aligned}$$

Hence from (3.50)

$$\begin{aligned} \frac{y(t_0)}{y'(t_0)} &= \int_{T(\gamma)}^{t_0} \frac{y'(t)}{y'(t_0)}\mathrm{d}t \\ &= t_0 - T(\gamma) + \int_{T(\gamma)}^{t_0}\left[(1+\frac{f(0)e^{-t}}{y'(t_0)})^{\frac{1}{n-1}}-1\right]\mathrm{d}t \\ &\quad +\mathcal{O}\left(\frac{(\log(g'))^{\beta+1}}{(g')^{\beta}}e^{-(g-\frac{n-1}{n}\gamma g')}+\frac{(\log(g'))}{(g')^3}\right). \end{aligned} \tag{3.53}$$

Let

$$A(\gamma) \overset{\text{def}}{=} \int_{T(\gamma)}^{t_0}\left[(1+\frac{f(0)e^{-t}}{y'(t_0)^{n-1}})^{\frac{1}{n-1}}-1\right]\mathrm{d}t.$$

If $f(0)=0$, then $A(\gamma)=0$. Let $f(0)\neq 0$ and $\theta_0 \overset{\text{def}}{=} -\log(f(0))+(n-1)\log(y'(t_0))$. Then,

$$\begin{aligned} A(\gamma) &= \int_{T(\gamma)+\theta_0}^{t_0+\theta_0}\left[(1+e^{-t})^{\frac{1}{n-1}}-1\right]\mathrm{d}t \quad \text{and} \\ t_0+\theta_0 &= (n+3)\log(g')-(n-1)\log(g')+\mathcal{O}(1)\to\infty \text{ as } \gamma\to\infty. \end{aligned}$$

Also from (3.41)

$$T(\gamma)+\theta_0 \geq (n-1)\log(g')-(n-1)\log(g')+\mathcal{O}(1)=\mathcal{O}(1).$$

Hence $A(\gamma)$ is bounded as $\gamma\to\infty$. Therefore we have from (3.53) and (3.50)

$$\begin{aligned} T(\gamma) &= t_0-\frac{y(t_0)}{y'(t_0)}+A(\gamma)+\mathcal{O}\left(\frac{(\log(g'))^{\beta+1}}{(g')^{\beta}}e^{-(g-(\frac{n-1}{n})\gamma g')}+\frac{(\log(g'))}{(g')^3}\right) \\ &= (g-\left(\frac{n-1}{n}\right)\gamma g')+(n-1)\log(\left(\frac{n-1}{n}\right)g')+\frac{(n-1)\alpha\gamma g''}{g'} \\ &\quad +A(\gamma)+\mathcal{O}\left(\frac{(\log(g'))^2}{g'}+\frac{(\log(g'))^{\beta+1}}{(g')^{\beta}}e^{-(g-(\frac{n-1}{n})\gamma g')}\right). \end{aligned}$$

This proves (3.47). Let $g-\left(\frac{n-1}{n}\right)\gamma g')\to\infty$ as $\gamma\to\infty$. Then for $t\in[T(\gamma),t_0]$ and from (3.46)

$$y'(t)^{n-1}=y'(t_0)^{n-1}+\int_t^{t_0}f(y(s))e^{-s}\mathrm{d}s=y'(t_0)^{n-1}+\mathcal{O}\left(e^{-T(\gamma)}\right).$$

Thus,

$$\begin{aligned} y'(t) &= y'(t_0)+\mathcal{O}\left(\frac{e^{-(g-(\frac{n-1}{n})\gamma g')}}{(g')^{n-1}}\right) \\ &= y'(\tilde{T})+\mathcal{O}\left(\frac{1}{(g')^3}+\frac{e^{-(g-(\frac{n-1}{n})\gamma g')}}{(g')^{n-1}}\right). \end{aligned}$$

This proves (3.48). From (3.52) we have

$$\begin{aligned} y'(T(\gamma)) &= y'(t_0)+y'(t_0)\left[(1+\frac{f(0)e^{-T(\gamma)}}{y'(t_0)^{n-1}})^{\frac{1}{n-1}}-1\right] \\ &\quad +\mathcal{O}\left(\frac{(\log(g'))^{\beta}}{(g')^{\beta}}e^{-(g-(\frac{n-1}{n})\gamma g')}+\frac{1}{(g')^3}\right) \end{aligned}$$

and (3.49) follows from (3.46). This proves the lemma. ☐

Proof of Theorem 2.2. : This follows from Lemma 3.9. ☐

4. Proof of Theorem 2.3.

First we look at the linearization of z which plays an important role in proving the theorem. Let $\theta \in \mathbb{R}$ and define $z_\theta(t) \overset{\text{def}}{=} z(t - \theta g') + \theta$, then z_θ satisfies

$$\left\{ \begin{array}{l} -((z_\theta')^{n-1})' = e^{g+g'(z_\theta-\gamma)-t} \\ z_\theta(\infty) = \gamma + \theta, \quad z_\theta'(\infty) = 0. \end{array} \right. \tag{4.1}$$

Let $V_2(t) = \frac{\partial z_\theta}{\partial \theta}|_{\theta=0}$. Then V_2 satisfies

$$V_2(t) = 1 - g'z'(t). \tag{4.2}$$

Differentiating (4.1) with respect to θ and evaluate at $\theta = 0$ to obtain

$$\left\{ \begin{array}{l} -((z')^{n-2}V_2')' = \dfrac{g'}{n-1}V_2 e^{g+g'(z-\gamma)-t} \\ V_2(\infty) = 1, \quad V_2'(\infty) = 0. \end{array} \right. \tag{4.3}$$

Moreover from (4.2), V_2 satisfies

$$\begin{aligned} V_2(t) &= \frac{1 - \frac{g'}{n}e^{\frac{g-t}{n-1}}}{1 + \left(\frac{n-1}{n}\right) g' e^{\frac{g-t}{n-1}}} = \frac{1 - \frac{1}{n-1}e^{\frac{T_1-t}{n-1}}}{1 + e^{\frac{T_1-t}{n-1}}} \\ &= -\frac{1}{n-1} + \left(\frac{n}{n-1}\right)\frac{1}{1+e^{\frac{T_1-t}{n-1}}}, \end{aligned} \tag{4.4}$$

$$V_2'(t) = \frac{n}{(n-1)^2}\frac{e^{\frac{T_1-t}{n-1}}}{(1+e^{\frac{T_1-t}{n-1}})^2}, \tag{4.5}$$

$$V_2''(t) = \frac{n}{(n-1)^3}\frac{e^{\frac{T_1-t}{n-1}}}{(1+e^{\frac{T_1-t}{n-1}})^2}\left[1 - \frac{2}{1+e^{\frac{T_1-t}{n-1}}}\right]. \tag{4.6}$$

Clearly $V_2' > 0$ and V_2 has exactly one zero at $t = S_0$ given by

$$S_0 = g + (n-1)\log\left(\frac{g'}{n}\right) = T_1 - (n-1)\log(n-1). \tag{4.7}$$

It is easy to see that

$$V_2(-\infty) = -\frac{1}{n-1}. \tag{4.8}$$

Let V_1 be the linearization of $y(t) = y(t, \gamma)$ as in section 2 (see (2.12)) and set

$$\begin{aligned} \delta &= \mathcal{O}(\log(g')), \quad T_\delta = T_1 - (n-1)\delta, \\ \rho_1(t) &\overset{\text{def}}{=} \frac{1}{(n-1)}f'(y(t)) = \frac{1}{n-1}e^{g(y(t))}g'(y(t)), \\ \rho_2(t) &\overset{\text{def}}{=} \frac{g'}{(n-1)}e^{g+g'(z(t)-\gamma)}. \end{aligned} \tag{4.9}$$

Then, V_1 and V_2 satisfy respectively:

$$-((y')^{n-2}V_1')' = \rho_1 V_1 e^{-t}, \tag{4.10}$$

$$-((z')^{n-2}V_2')' = \rho_2 V_2 e^{-t}. \tag{4.11}$$

For $T(\gamma) \leq t < \eta$, integrating (4.10) and (4.11) gives

$$V_1'(t) = \left(\frac{y'(\eta)}{y'(t)}\right)^{n-2} V_1'(\eta) + \frac{1}{y'(t)^{n-2}} \int_t^\eta \rho_1 V_1 e^{-s} \mathrm{d}s. \tag{4.12}$$

Thus,

$$V_1(t) = V_1(\eta) - V_1'(\eta) \int_t^\eta \left(\frac{y'(\eta)}{y'(s)}\right)^{n-2} \mathrm{d}s - \int_t^\eta \rho_1 V_1 e^{-s} \left(\int_t^s \frac{\mathrm{d}\theta}{y'(\theta)^{n-2}}\right) \mathrm{d}s. \tag{4.13}$$

Similarly,

$$V_2'(t) = \left(\frac{z'(\eta)}{z'(t)}\right)^{n-2} V_2'(\eta) + \frac{1}{z'(t)^{n-2}} \int_t^\eta \rho_2 V_2 e^{-s} \mathrm{d}s \tag{4.14}$$

and

$$V_2(t) = V_2(\eta) - V_2'(\eta) \int_t^\eta \left(\frac{z'(\eta)}{z'(s)}\right)^{n-2} \mathrm{d}s - \int_t^\eta \rho_2 V_2 e^{-s} \left(\int_t^s \frac{\mathrm{d}\theta}{z'(\theta)^{n-2}}\right) \mathrm{d}s. \tag{4.15}$$

Let

$$\begin{aligned} \rho_3(t) &\overset{\text{def}}{=} \big((y'(t)^{n-2} - z'(t)^{n-2})V_2'(t)\big)' \\ &= (n-2)(y'(t)^{n-3}y''(t) - z'(t)^{n-3}z''(t))V_2' + (y'(t)^{n-2} - z'(t)^{n-2})V_2''. \end{aligned} \tag{4.16}$$

Then we have

$$\begin{aligned} \big((y')^{n-2}(V_1'V_2 - V_1V_2')\big)' &= ((y')^{n-2}V_1')'V_2 - ((y')^{n-2}V_2')'V_1 \\ &= -\rho_1 V_1 V_2 e^{-t} + \rho_2 V_1 V_2 e^{-t} - \rho_3(t) V_1. \end{aligned}$$

Therefore,

$$\begin{aligned} V_1'(t)V_2(t) - V_1(t)V_2'(t) &= \left(\frac{y'(\eta)}{y'(t)}\right)^{n-2} (V_1'(\eta)V_2(\eta) - V_1(\eta)V_2'(\eta)) + \\ &\quad + \frac{1}{y'(t)^{n-2}} \int_t^\eta (\rho_1 - \rho_2) V_1 V_2 e^{-s} \mathrm{d}s + \frac{1}{y'(t)^{n-2}} \int_t^\eta \rho_3 V_1(s) \mathrm{d}s. \end{aligned} \tag{4.17}$$

In all the future lemmas, the statement deals for large γ; that is there exists $\gamma_0 > 0$ such that for $\gamma > \gamma_0$, the lemma is true. Note that γ_0 may be different for different lemmas. With abuse of notation, we delete the statement $\gamma > \gamma_0$ or for large γ.

We need the following estimates proved in Section 3 (see (3.16) and (3.18)) and some more estimates. Let $t \geq T_\delta$, then

$$g(y(t)) = g + g'(z(t) - \gamma) + \mathcal{O}\left(\frac{g''}{(g')^2}(\log(1 + e^\delta))^2\right), \tag{4.18}$$

$$g'(y(t)) = g' - \frac{ng''}{g'} \log(1 + e^\delta) + \mathcal{O}\left(\frac{g''}{(g')^2}(\log(1 + e^\delta))^2\right) \tag{4.19}$$

$$\begin{aligned} g''(y(t)) &= g'' + \mathcal{O}\left(\frac{g'''}{g'}\log(1+e^\delta)\right) = g''\left[1+\mathcal{O}\left(\frac{1}{\gamma g'}\log(1+e^\delta)\right)\right] \\ &= g''\left[1+\mathcal{O}\left(\frac{\log(1+e^\delta)}{(g')^{1+\frac{1}{q-1}}}\right)\right] = g''\left[1+\mathcal{O}\left(\frac{\log(1+e^\delta)}{(g')^n}\right)\right]. \end{aligned} \tag{4.20}$$

Since $g' \sim \gamma^{q-1}$, $1 < q \le \dfrac{n}{n-1}$, $\dfrac{g''}{(g')^2} \sim \left(\dfrac{\gamma g''}{g'}\right)\dfrac{1}{\gamma g'} \sim \dfrac{1}{(g')^{\frac{q}{q-1}}}$ and $n \le \dfrac{q}{q-1}$. Hence,

$$\begin{aligned} \frac{g''(y(t))}{g'(y(t))} &= \frac{g''}{g'}\left[1+\mathcal{O}\left(\frac{\log(1+e^\delta)}{(g')^n}\right)\right]\left[1+\frac{ng''}{(g')^2}\log(1+e^\delta)+\mathcal{O}\left(\frac{g''}{(g')^3}(\log(1+e^\delta))^2\right)\right] \\ &= \frac{g''}{g'}\left[1+\mathcal{O}\left(\frac{g''\log(1+e^\delta)}{(g')^2}\right)\right] \end{aligned} \tag{4.21}$$

and

$$\frac{g'''(y(t))}{g'(y(t))} = \mathcal{O}\left(\frac{g'''}{g'}\right) = O\left(\frac{\gamma g'''}{g''}\frac{g''}{\gamma g'}\right) = \mathcal{O}\left(\frac{g''}{(g')^n}\right). \tag{4.22}$$

Therefore,

$$\begin{aligned} \rho_1(t) &= \frac{1}{n-1}g'(y(t))e^{g(y(t))} \\ &= \frac{g'}{(n-1)}\left(1+\mathcal{O}\left(\frac{g''}{(g')^2}\log(1+e^\delta)\right)\right)e^{g+g'(z(t)-\gamma)+\mathcal{O}\left(\frac{g''}{(g')^2}(\log(1+e^\delta))^2\right)}, \end{aligned}$$

and

$$\rho_1(t) = \rho_2(t)e^{\mathcal{O}\left(\frac{g''}{(g')^2}\log(1+e^\delta)^2\right)}. \tag{4.23}$$

Let $T_\delta \le t_1 < t_2 < \infty$, then from (3.7) and by integration by parts

$$\begin{aligned} 0 < \int_{t_1}^{t_2}\rho_2 e^{-s}\left(\int_{t_1}^{s}\frac{\mathrm{d}\theta}{(z'(\theta))^{n-2}}\right)\mathrm{d}s &= \frac{g'}{(n-1)}\int_{t_1}^{t_2}e^{g+g'(z-\gamma)-s}\left(\int_{t_1}^{s}\frac{\mathrm{d}\theta}{(z'(\theta))^{n-2}}\right)\mathrm{d}s \\ &= \frac{g'}{n-1}\left[z(t_2)-z(t_1)-z'(t_2)\int_{t_1}^{t_2}\left(\frac{z'(t_2)}{z'(t)}\right)^{n-2}\mathrm{d}t\right] \\ &\le \frac{g'}{n-1}|z(t_2)-z(t_1)| \le \frac{n}{(n-1)^2}|t_2-t_1|. \end{aligned} \tag{4.24}$$

Setting $t_2 = \infty$, we get

$$\begin{aligned} 0 < \int_t^\infty(\rho_2 e^{-s})\left(\int_t^s\frac{\mathrm{d}\theta}{(z'(\theta))^{n-2}}\right)\mathrm{d}s &\le \frac{g'}{n-1}(\gamma - z(t)) \\ &\le \frac{n}{n-1}\log(1+e^{\frac{T_1-t}{n-1}}) \end{aligned} \tag{4.25}$$

and

$$\frac{1}{z'(t_1)^{n-2}}\int_{t_1}^{t_2}\rho_2 e^{-s}\mathrm{d}s = \frac{g'}{(n-1)}\left(z'(t_1)-\left(\frac{z'(t_2)}{z'(t_1)}\right)^{n-2}z'(t_2)\right) \le \frac{n}{(n-1)^2}. \tag{4.26}$$

Idea of the proof: Basic idea is

(1) Let $S < S_1 < \infty$ be the first turning point and first zero of V_1 before infinity. Then in the first few lemmas, we show that S_1 exists. Then estimates about $|V_1(t)-V_2(t)|+|V_1'(t)-V_2'(t)|$ for $t \in [S,\infty)$ are established.

(2) Since $V_2(t) \to -\dfrac{1}{n-1}$ as $t \to -\infty$, hence for $T(\gamma) < t < S$, V_1 and V_2 depart. Therefore we need a better estimate of S and this is achieved by using the identity (4.61) in Lemma 4.6. This identity plays an important role in proving the crucial estimate (4.66) in Lemma 4.9 which yields the necessary result.



Lemma 4.1. *Let* $T_\delta \le t \le \eta \le \infty$, $A = \mathcal{O}\left(\dfrac{g''}{(g')^2}(\log(1+e^\delta))^2\right)$ *and*

$$M_\eta(t) = \max\left\{|V_1(s) - V_2(s)|;\ t \le s \le \eta\right\}. \tag{4.27}$$

Then for γ *large,*

$$\begin{aligned} M_\eta(t) &\le |V_1(\eta) - V_2(\eta)| + 2|\eta - t||V_1'(\eta) - V_2'(\eta)| + \frac{2n}{n-1}\log(1+e^\delta)M_\eta(t) \\ &+ 2A\left[|V_2'(\eta)|(\eta - t) + \log(1 + e^{\frac{T_1 - t}{n-1}})\right], \end{aligned} \tag{4.28}$$

$$M_\infty(t) \le 4\log(1 + e^{\frac{T_1-t}{n-1}})[A + M_\infty(t)] \ \textit{and} \tag{4.29}$$

$$|V_1'(t) - V_2'(t)| \le 2|V_1'(\eta) - V_2'(\eta)| + 2A(|V_2'(\eta)| + V_2(\eta)) + 4M_\eta(t). \tag{4.30}$$

Proof. From (3.17), (4.23) and (4.15), we have for $T_\delta \le t \le s \le \eta \le \infty$,

$$\begin{aligned} V_1(s) &= V_1(\eta) - V_1'(\eta)\int_s^\eta \left(\frac{y'(\eta)}{y'(\theta)}\right)^{n-2} \mathrm{d}\theta - \int_s^\eta \rho_1 V_1 e^{-\theta}\left(\int_s^\theta \frac{\mathrm{d}x}{y'(x)^{n-2}}\right)\mathrm{d}\theta \\ &= V_1(\eta) - V_1'(\eta)e^A\int_s^\eta \left(\frac{z'(\eta)}{z'(\theta)}\right)^{n-2}\mathrm{d}\theta - e^A\int_s^\eta \rho_2 V_1 e^{-\theta}(\int_s^\theta \frac{\mathrm{d}x}{z'(x)^{n-2}})\mathrm{d}\theta \\ &= V_1(\eta) - (V_1'(\eta) - V_2'(\eta))e^A\int_s^\eta \left(\frac{z'(\eta)}{z'(\theta)}\right)^{n-2}\mathrm{d}\theta - V_2'(\eta)e^A\int_s^\eta \left(\frac{z'(\eta)}{z'(\theta)}\right)^{n-2}\mathrm{d}\theta \\ &\quad -e^A\int_s^\eta \rho_2 V_2 e^{-\theta}(\int_s^\theta \frac{\mathrm{d}x}{z'(x)^{n-2}})\mathrm{d}\theta + e^A\int_s^\eta \rho_2(V_2 - V_1)e^{-\theta}(\int_s^\theta \frac{\mathrm{d}x}{z'(x)^{n-2}})\mathrm{d}\theta. \end{aligned}$$

Hence

$$\begin{aligned} V_1(s) - V_2(s) &= (V_1(\eta) - V_2(\eta)) - (V_1'(\eta) - V_2'(\eta))e^A \int_s^\eta \left(\frac{z'(\eta)}{z'(\theta)}\right)^{n-2} \mathrm{d}\theta \\ &\quad +(e^A - 1)\left[-V_2'(\eta)\int_s^\eta \left(\frac{z'(\eta)}{z'(\theta)}\right)^{n-2} \mathrm{d}\theta - \int_s^\eta \rho_2 V_2 e^{-\theta}\Big(\int_s^\theta \frac{\mathrm{d}x}{(z'(x))^{n-2}}\Big)\mathrm{d}\theta\right] \\ &\quad +e^A \int_s^\eta \rho_2(V_2 - V_1)e^{-\theta}\Big(\int_s^\theta \frac{\mathrm{d}x}{z'(x)^{n-2}}\Big)\mathrm{d}\theta. \end{aligned} \tag{4.31}$$

Since $z'(\eta) \le z'(\theta)$ for $\theta \le \eta$ and $0 \le V_2 \le 1$, we have by using (4.25) for $\eta \ne \infty$

$$\begin{aligned} |V_1(s) - V_2(s)| &\le |V_1(\eta) - V_2(\eta)| + e^A(\eta - s)|V_1'(\eta) - V_2'(\eta)| \\ &\quad +(e^A - 1)\left(|V_2'(\eta)|(\eta - s) + \left(\frac{n}{n-1}\right)\log(1 + e^{\frac{T_1 - s}{n-1}})\right) \\ &\quad + \left(\frac{n}{n-1}\right) e^A \log(1 + e^{\frac{T_1 - s}{n-1}})M_\eta(s). \end{aligned} \tag{4.32}$$

If $\eta = \infty$, using (2.3) and (2.12), we get

$$|V_1(s) - V_2(s)| \le (e^A - 1)\frac{n}{n-1}\log\left(1 + e^{\frac{T_1 - s}{n-1}}\right) + \frac{n}{n-1}e^A \log\left(1 + e^{\frac{T_1 - s}{n-1}}\right) M_\infty(t).$$

For γ large, we have $e^A \le 1 + 2A$ and $e^A \le 2$, hence by taking the maximum of $|V_1 - V_2|$ in $[t, \eta]$ we obtain if $\eta < \infty$

$$\begin{aligned} M_\eta(t) &\le |V_1(\eta) - V_2(\eta)| + 2|t - \eta||V_1'(\eta) - V_2'(\eta)| \\ &\quad +2A\left(|V_2'(\eta)||\eta - t| + \frac{n}{n-1}\log\left(1 + e^{\frac{T_1 - t}{n-1}}\right)\right) \\ &\quad +4\log\left(1 + e^{\frac{T_1 - t}{n-1}}\right) M_\eta(t), \end{aligned}$$

and if $\eta = \infty$, the estimate

$$M_\infty(t) \le 4\log(1 + e^{\frac{T_1 - t}{n-1}})(A + M_\infty(t)).$$

This proves (4.28) and (4.29). Next,

$$\begin{aligned} V_1'(t) &= \left(\frac{y'(\eta)}{y'(t)}\right)^{n-2} V_1'(\eta) + \frac{1}{y'(t)^{n-2}}\int_t^\eta \rho_1 V_1 e^{-s}\mathrm{d}s \\ &= e^A V_1'(\eta)\left(\frac{z'(\eta)}{z'(t)}\right)^{n-2} + \frac{e^A}{z'(t)^{n-2}}\int_t^\eta \rho_2(V_1 - V_2)e^{-s}\mathrm{d}s + \frac{e^A}{z'(t)^{n-2}}\int_t^\eta \rho_2 V_2 e^{-s}\mathrm{d}s \\ &= e^A\left(V_2'(\eta)\left(\frac{z'(\eta)}{z'(t)}\right)^{n-2} + \frac{1}{z'(t)^{n-2}}\int_t^\eta \rho_2 V_2 e^{-s}\mathrm{d}s\right) \\ &\quad +e^A(V_1'(\eta) - V_2'(\eta))\left(\frac{z'(\eta)}{z'(t)}\right)^{n-2} + \frac{e^A}{z'(t)^{n-2}}\int_t^\eta \rho_2(V_1 - V_2)e^{-s}\mathrm{d}s. \end{aligned}$$

Hence from (4.14), (4.26), we have

$$|V_1'(t) - V_2'(t)| \le 2|V_1'(\eta) - V_2'(\eta)| + 2A(|V_2'(\eta)| + V_2(\eta)) + 4M_\eta(t).$$

This proves the lemma. □

In the next few lemmas we estimate the first zero S_1 and the first turning point S defined by

$$S_1 \overset{\text{def}}{=} \inf\{t \,:\, V_1(s) > 0, \quad \forall s \in (t, \infty)\}, \tag{4.33}$$

$$S \overset{\text{def}}{=} \inf\{t : V_1'(s) > 0, \quad \forall s \in (t, \infty)\}, \tag{4.34}$$

Lemma 4.2. *Let S_0 be the zero of V_2 as defined in (4.7). Then there exists a $k_1 > 0$, $C_3 > 0$ and $\gamma_0 > 0$ such that for all $\gamma \geq \gamma_0$, we have*

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^4}\right) \quad \forall t \geq T_1 + 2(n-1)\log(g'), \tag{4.35}$$

$$S_1 - S_0 = \mathcal{O}\left(\frac{g''}{(g')^2}\right), \tag{4.36}$$

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right) \quad \forall t \geq S_1 - (n-1)k_1, \tag{4.37}$$

$$V_2(S_1 - (n-1)k_1) \leq -C_3. \tag{4.38}$$

Proof. Let $\delta = -2\log(g')$, then $A = \mathcal{O}\left(\frac{g''}{(g')^2}(\log(1+e^\delta))^2\right) = \mathcal{O}\left(\frac{g''}{(g')^4}\right)$ and hence for $t \geq T_\delta = T_1 - (n-1)\delta$ and using Lemma 4.1, we have

$$|V_1(t) - V_2(t)| = \mathcal{O}\left(\frac{g''}{(g')^4}\right).$$

Let $\eta = \infty$ in (4.30) and $t \geq T_1 - (n-1)\delta$, we have

$$|V_1'(t) - V_2'(t)| = 2A + M_\infty(t) = \mathcal{O}\left(\frac{g''}{(g')^4}\right).$$

This proves (4.35). Let $k_0 > 0$ be large such that $4\log(1+e^{-k_0}) < \frac{1}{2}$ and let $-\delta = k_0$, then $A = \mathcal{O}\left(\frac{g''}{(g')^2}\right)$. Then from (4.29) and for $t \geq T_1 + (n-1)k_0$

$$\frac{M_\infty(t)}{2} \leq 4\log(1 + e^{\frac{T_1 - t}{n-1}}))A.$$

Hence $M_\infty(t) = \mathcal{O}(A) = \mathcal{O}\left(\frac{g''}{(g')^2}\right)$. From (4.30) taking $\eta = \infty$ we obtain for $t \geq T_1 + (n-1)k_0$ the estimate $|V_1'(t) - V_2'(t)| = \mathcal{O}(A)$. Hence for $t \geq T_1 + (n-1)k_0$, we have

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right). \tag{4.39}$$

Claim 1. $S_1 < T_1 + (n-1)k_0$.

Proof of Claim 1: Since $T_1 + (n-1)k_0 = g' + (n-1)\log\left(\frac{g'(n-1)}{n}\right) + (n-1)k_0 = S_0 + (n-1)k_0 + (n-1)\log(n-1)$, hence if $S_1 \geq T_1 + (n-1)k_0$, then $S_1 > S_0$ and $S_0 - S_1 < -(n-1)(k_0 + \log(n-1)) \leq -(n-1)k_0$. This along with (4.39), (4.4) gives

$$\frac{1 - e^{\frac{S_0 - S_1}{n-1}}}{1 + e^{\frac{T_1 - S_1}{n-1}}} = V_2(S_1) = V_2(S_1) - V_1(S_1) = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

Thus,

$$1 = e^{\frac{S_0 - S_1}{n-1}} + \mathcal{O}\left((1 + e^{\frac{T_1 - S_1}{n-1}})\frac{g''}{(g')^2}\right) \leq e^{-k_0} + \mathcal{O}\left((1 + e^{-k_0})\frac{g''}{(g')^2}\right) < 1$$

which is a contradiction and the claim follows. ☐

Claim 2. There exists a $C_1 > 0$ such that

$$S_1 \geq T_1 + (n-1)k_0 - C_1. \tag{4.40}$$

Proof of Claim 2: Let $S_2 = T_1 + (n-1)k_0$, then from (3.27), for $T(\gamma) \leq t \leq S_2$, we have $y'(S_2) \geq z'(S_2)$ and $y'(t) \leq y'(T(\gamma))$. Hence from (3.31), there exists a $C_0 > 0$ such that for γ large

$$\frac{y'(S_2)}{y'(t)} \geq \frac{z'(S_2)}{y'(T(\gamma))} = \frac{\frac{n}{g'(n-1)}(\frac{e^{-k_0}}{1+e^{-k_0}})}{(1+C_0)(\frac{n}{g'(n-1)} + \mathcal{O}\left(\frac{1}{(g')^3}\right))} \geq C_2$$

where $C_2 > 0$ independent of γ (C_2 depends on γ_0 but is independent of $\gamma > \gamma_0$). Hence from Claim 1, (2.12) and (4.39)

$$\begin{aligned} V_2(S_2) + O\left(\frac{g''}{(g')^2}\right) &= V_1(S_2) = V_1(S_2) - V_1(S_1) \\ &\geq V_1'(S_2)\int_{S_1}^{S_2}\left(\frac{y'(S_2)}{y'(t)}\right)^{n-2}\mathrm{d}t \\ &\geq C_2V_1'(S_2)(S_2 - S_1) = C_2(V_2'(S_2) + \mathcal{O}\left(\frac{g''}{(g')^2}\right))(S_2 - S_1). \end{aligned}$$

Since $V_2(S_2) = \dfrac{1 - \frac{e^{-k_0}}{n-1}}{1+e^{-k_0}}$, $V_2'(S_2) = \dfrac{n}{(n-1)^2}\dfrac{e^{-k_0}}{1+e^{-k_0}}$, hence there exists a $C_1 > 0$ independent of γ such that $S_2 - S_1 \leq C_1$. This proves the claim. □

Claim 3. For $t \geq S_1$

$$S_1 = S_0 + \mathcal{O}\left(\frac{g''}{(g')^2}\right), \tag{4.41}$$

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right). \tag{4.42}$$

Proof of Claim 3: From Claims 1 and 2,we see that $S_1 = T_1 + (n-1)\delta_0$ and $\delta_0 = \mathcal{O}(1)$. Hence for $t \in [S_1, \infty)$,we have $0 \leq V_1(t) \leq 1$, $S_2 - S_1 = \mathcal{O}(1)$ and from (4.23) $\rho_1 = \rho_2 e^{\mathcal{O}\left(\frac{g''}{(g')^2}\right)}$. Now, using (4.26) and (3.17) we have

$$\begin{aligned} \frac{1}{y'(S_1)^{n-2}}\int_{S_1}^{S_2}(\rho_1 - \rho_2)V_1V_2e^{-t}\mathrm{d}t &= \mathcal{O}\left(\frac{g''}{(g')^2}\right)\frac{1}{z'(S_1)^{n-2}}\int_{S_1}^{S_2}\rho_2e^{-t}\mathrm{d}t \\ &= \mathcal{O}\left(\frac{g''}{(g')^2}\right), \end{aligned} \tag{4.43}$$

$$V_1'(t) = \frac{1}{y'(t)^{n-2}}\int_t^\infty \rho_1e^{-s}V_1\mathrm{d}s = \mathcal{O}\left(\frac{1}{z'(t)^{n-2}}\int_t^\infty \rho_2e^{-s}\mathrm{d}s\right) = \mathcal{O}(1), \tag{4.44}$$

$$\begin{aligned} \frac{1}{y'(S_1)^{n-2}}\left(y'(t)^{n-2} - z'(t)^{n-2}\right)V_2'(t)V_1(t) &= \frac{V_1(t)}{y'(S_1)^{n-2}}\int_t^\infty(\rho_1 - \rho_2)e^{-t}\mathrm{d}t + \mathcal{O}\left(\frac{g''}{(g')^2}\right) \\ &= \mathcal{O}\left(\frac{g''}{(g')^2}\right) \end{aligned}$$

and similarly

$$\frac{1}{y'(S_1)^{n-2}}\int_{S_1}^{S_2}(y'(t)^{n-2} - z'(t)^{n-2})V_2'(t)V_1(t)\mathrm{d}t = \mathcal{O}\left(\frac{g''}{(g')^2}(S_2 - S_1)\right) = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

Therefore from the above estimates, we have

$$\begin{aligned}\frac{1}{y'(S_1)^{n-2}}\int_{S_1}^{S_2}\rho_3 V_1 \mathrm{d}t &= \frac{1}{y'(S_1)^{n-2}}\left[(y'(t)^{n-2}-z'(t)^{n-2})V_2'(t)V_1(t)\right]_{S_1}^{S_2} \\ &\quad -\frac{1}{y'(S_1)^{n-2}}\int_{S_1}^{S_2}(y'(t)^{n-2}-z'(t)^{n-2})V_2'(t)V_1'(t)\mathrm{d}t \\ &= \mathcal{O}\left(\frac{g''}{(g')^2}\right).\end{aligned} \tag{4.45}$$

Taking $\eta = S_2$, $t = S_1$ in (4.17) and from the above estimates we have,

$$\begin{aligned}V_1'(S_1)V_2(S_1) &= \left(\frac{y'(S_2)}{y'(S_1)}\right)^{n-2}\left[(V_1'(S_2)-V_2'(S_2))V_2(S_2)+(V_2(S_2)-V_1(S_2))V_1'(S_2)\right]+\mathcal{O}\left(\frac{g''}{(g')^2}\right) \\ &= \mathcal{O}\left(\frac{g''}{(g')^2}\right).\end{aligned}$$

From Claim 1, we have

$$V_1'(S_1) \geq V_1'(S_2)\frac{(y')^{n-2}(S_2)}{(y')^{n-2}(S_1)} = V_2'(S_2)\mathcal{O}\left(e^{\frac{g''}{(g')^4}}\right)+\mathcal{O}\left(\frac{g''}{(g')^2}\right) \geq C_3 > 0$$

for some C_3 independent of γ. Hence $V_2(S_1) = \mathcal{O}\left(\frac{g''}{(g')^2}\right)$ and

$$\left|\frac{1-e^{\frac{S_0-S_1}{n-1}}}{1+e^{\frac{T_1-S_1}{n-1}}}\right| = |V_2(S_1)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

Since $T_1 - S_1 = \mathcal{O}(1)$ and hence from the above equation,we get $S_0 - S_1 = \mathcal{O}\left(\frac{g''}{(g')^2}\right)$. This proves (4.40). Also

$$|V_2(S_1) - V_1(S_1)| = |V_2(S_1)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right). \tag{4.46}$$

We now have two situations to handle, but before that we shall prove a few other estimates. Let $\varepsilon > 0$ such that $\frac{n}{(n-1)}\varepsilon = \frac{1}{2}$ and taking $t = S_0$ and $\eta = S_0 + \varepsilon$ in (4.32), we get on $[S_0, S_0+\varepsilon]$

$$\begin{aligned}|V_1(s)-V_2(s)| &\leq |V_1(\eta)-V_2(\eta)| + e^A|V_1'(\eta)-V_2'(\eta)||s-\eta| \\ &+ (e^A-1)\left[|V_2'(\eta)||s-\eta| + \frac{n}{n-1}\log(1+e^{\frac{T_1-S_0}{n-1}})\right] + \frac{n}{n-1}|s-\eta|M_\eta(s)\end{aligned}$$

which now easily gives the estimate

$$M_\eta(s) \leq C_0\left(|V_1(\eta)-V_2(\eta)| + |V_1'(\eta)-V_2'(\eta)| + \mathcal{O}\left(\frac{g''}{(g')^2}\right)\right) \tag{4.47}$$

for all $s \in [S_0, S_0+\varepsilon]$ and some constant $C_0 > 0$ independent of γ.

Since $V_2(S_0+\varepsilon) = \dfrac{1-e^{\varepsilon}}{1+e^{\frac{T_1-S_0}{n-1}}} \geq C_1 > 0$ depending only on ε, observe that if we replace S_1 by S_0 in (4.43), (4.44) and (4.45), still all the estimates hold. Hence for $t \in [S_0+\varepsilon, S_2]$ and from previous claims,

$$\begin{aligned}
V_1'(t)V_2(t) - V_1(t)V_2'(t) &= \left(\frac{y'(S_2)}{y'(t)}\right)^{n-2} (V_1'(S_2)V_2(S_2) - V_1(S_2)V_2'(S_2)) + \mathcal{O}\left(\frac{g''}{(g')^2}\right) \\
&= \left(\frac{y'(S_2)}{y'(t)}\right)^{n-2} (V_1'(S_2) - V_2'(S_2))V_2(S_2) + \\
&\quad +(V_2(S_2) - V_1(S_2))V_2'(S_2) + \mathcal{O}\left(\frac{g''}{(g')^2}\right) \\
&= \mathcal{O}\left(\frac{g''}{(g')^2}\right).
\end{aligned} \tag{4.48}$$

Note that we needed $S_0+\varepsilon < S_2$, or in other words we needed $\dfrac{1}{2}\dfrac{(n-1)}{n} =: \varepsilon < (n-1)[k_0 + \log(n-1)]$. Choose $k_0 > 0$ large such that the required estimate holds.

Similar to Claim 1, we also have $S_0+\varepsilon > S_1$ and hence $V_1(S_0+\varepsilon) > 0$. In particular,

$$V_1(t) \geq C_6 > 0 \text{ for } t \in [S_0+\varepsilon, S_2]$$

Proof of $S_0+\varepsilon > S_1$. Suppose not, then we must have $S_0 - S_1 \leq -\varepsilon$. From the definition of V_2, we get

$$V_2(S_1) = \frac{1-e^{\frac{S_0-S_1}{n-1}}}{1+(n-1)e^{\frac{S_0-S_1}{n-1}}} = V_2(S_1) - V_1(S_1) = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

This implies for γ large enough

$$1 = e^{\frac{S_0-S_1}{n-1}} + \mathcal{O}\left(\left(1+e^{\frac{S_0-S_1}{n-1}}\right)\frac{g''}{(g')^2}\right) \leq e^{-\varepsilon} + \mathcal{O}\left([1+(n-1)e^{-\varepsilon}]\frac{g''}{(g')^2}\right) < 1$$

which gives the necessary contradiction and this proves that $S_0+\varepsilon > S_1$. □

As a consequence of $S_0+\varepsilon > S_1$, we see that $V_1(t) \geq C_6 > 0$ for $t \in [S_0+\varepsilon, S_2]$. Since $V_2(S_0) = 0$, we also have $V_2(s) \geq \tilde{C}_6 > 0$ for $s \in [S_0+\varepsilon, S_2]$. This allows us to divide by V_1 and V_2 in (4.48) to get

$$\frac{V_1(t)}{V_2(t)} = \frac{V_1(S_2)}{V_2(S_2)} e^{\mathcal{O}\left(\frac{g''}{(g')^2}\right)}.$$

Integrating from t to S_2, we now get

$$\begin{aligned}
V_1(t) - V_2(t) &= \frac{V_1(S_2)}{V_2(S_2)}V_2(t) - V_2(t) + \mathcal{O}\left(\frac{g''}{(g')^2}\right) \\
&= (V_1(S_2) - V_2(S_2))\frac{V_2(t)}{V_2(S_2)} + \mathcal{O}\left(\frac{g''}{(g')^2}\right) = \mathcal{O}\left(\frac{g''}{(g')^2}\right).
\end{aligned}$$

Hence from (4.30) and (4.39), $|V_1'(t) - V_2'(t)| = \mathcal{O}\left(\dfrac{g''}{(g')^2}\right)$. That is for all $t \in [S_0+\varepsilon, S_2]$,

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

Using this in (4.47), we get

$$|V_1(t) - V_2(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right) \text{ for } t \in [S_0, S_0+\varepsilon]$$

and now making use of (4.30), we get

$$|V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right) \text{ for } t \in [S_0, S_0+\varepsilon].$$

Let

$$I = \begin{cases} [S_0, S_1] & \text{if } S_0 < S_1 \\ [S_1, S_0] & \text{if } S_0 \geq S_1. \end{cases}$$

Case i: $I = [S_0, S_1]$. In this case, our previous calculations automatically gives the desired estimate

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right) \text{ for } t \in I$$

Case ii: $I = [S_1, S_0]$. From (4.46), we see that for $t \in I$,

$$\begin{aligned} V_1(t) - V_2(t) &= V_1(S_1) - V_2(S_1) - \int_t^{S_1} (V_1'(\theta) - V_2'(\theta))\mathrm{d}\theta \\ &= \mathcal{O}\left(\frac{g''}{(g')^2}\right) + \mathcal{O}(|S_1 - S_0|) = \mathcal{O}\left(\frac{g''}{(g')^2}\right). \end{aligned} \tag{4.49}$$

Hence from (4.28), (4.30), we obtain

$$|V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{g''}{(g')^2}\right) \text{ for } t \in I. \tag{4.50}$$

This completes the proof of Claim 3. □

Let $k_1 > 0$ to be chosen later and $S_3 = S_1 - (n-1)k_1$. Then

$$S_3 - S_0 = S_1 - S_0 - (n-1)k_1 = \mathcal{O}\left(\frac{g''}{(g')^2}\right) - (n-1)k_1$$

and

$$T_1 - S_3 = T_1 - S_0 - (S_1 - S_0) + (n-1)k_1 = \mathcal{O}(1).$$

Hence

$$V_2(S_3) = \frac{1 - e^{k_1 + O\left(\frac{g''}{(g')^2}\right)}}{1 + e^{O(1)}} \leq -C_3 \tag{4.51}$$

which proves (4.38).

Consider $s \in [S_3, S_1]$ and now using (4.31), we get with $\eta = S_1$

$$\begin{aligned} V_1(s) - V_2(s) \leq\ & |V_1(S_1) - V_2(S_1)| - |V_1'(S_1) - V_2'(S_1)|e^A \int_s^{S_1} \left(\frac{z'(S_1)}{z'(\theta)}\right)^{n-2} \mathrm{d}\theta \\ & + (e^A - 1)\left[|V_2'(S_1)| \int_s^{S_1} \left(\frac{z'(S_1)}{z'(\theta)}\right)^{n-2} \mathrm{d}\theta + \int_s^{S_1} \rho_2 V_2 e^{-\theta}\left(\int_s^{\theta} \frac{\mathrm{d}x}{(z'(x))^{n-2}}\right)\mathrm{d}\theta\right] \\ & + e^A \int_s^{S_1} \rho_2 |V_2 - V_1| e^{-\theta}\left(\int_s^{\theta} \frac{\mathrm{d}x}{z'(x)^{n-2}}\right)\mathrm{d}\theta \\ \leq\ & |V_1(S_1) - V_2(S_1)| - |V_1'(S_1) - V_2'(S_1)| 2(n-1)k_1 \\ & + (e^A - 1)\left[|V_2'(S_1)|(n-1)k_1 + \left(\frac{n-1}{n}\right) \log\left(1 + e^{\frac{T_1 - S_3}{n-1}}\right)\right] \\ & + e^A M_{S_3}(s) \frac{n}{(n-1)^2} k_1 (n-1). \end{aligned}$$

Now choose k_1 small such that $e^A\dfrac{n}{(n-1)}k_1 \leq \dfrac{1}{2}$ which implies

$$M_{S_3}(S_1) = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

This completes the proof of the Lemma. ☐

Lemma 4.3. *Let* $k_1 > 0$ *and* $S_3 = S_1 - (n-1)k_1$ *be as in Lemma 4.2. Let* $\delta > 0$, $\delta = \mathcal{O}(\log(g'))$ *and* $T_\delta = T_1 - (n-1)\delta$. *Define*

$$S_4 = \max\{T_\delta, S\}. \tag{4.52}$$

Then for all $t \in [S_4, S_3]$, *we have*

$$|V_1(t) - V_2(t)| + |V_1'(t) - V_2'(t)| = \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right). \tag{4.53}$$

Proof. From Lemma 4.2 and (4.31),

$$V_1(S_3) = V_2(S_3) + \mathcal{O}\left(\frac{g''}{(g')^2}\right) < -C_3 + \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

Hence for $t \in [S_4, S_3]$ and γ large,we have

$$V_1(S_3) \leq -\frac{C_3}{2} < 0, \quad |S_4 - S_3| = \mathcal{O}(\delta) \quad \log(1+e^\delta) = \mathcal{O}(\delta),$$
$$0 \leq V_1'(t) \leq \frac{y'(S_4)^{n-2}}{y'(t)^{n-2}} V_1'(S_3) = V_2'(S_3) + \mathcal{O}(1) = \mathcal{O}(1).$$

Since $V_1(t) < 0$ for $t \in [S_4, S_3]$, by using (4.23), we obtain

$$\begin{aligned} \frac{1}{y'(t)^{n-2}}\int_t^{S_3}(\rho_1 - \rho_2)V_1V_2e^{-s}\mathrm{d}s &= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}|\frac{1}{y'(t)^{n-2}}\int_t^{S_3}\rho_1V_1e^{-s}\mathrm{d}s|\right) \\ &= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}(V_1'(t) - \left(\frac{y'(S_3)}{y'(t)}\right)^{n-2}V_1'(S_3))\right) \\ &= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right). \end{aligned} \tag{4.54}$$

For $s \in [t, S_3]$ and from (3.17), we have

$$\frac{1}{y'(t)^{n-2}}\left[y'(s)^{n-2} - z'(s)^{n-2}\right] = \mathcal{O}\left(\left(\left(\frac{y'(s)}{z'(s)}\right)^{n-2} - 1\right)\left(\frac{z'(s)}{z'(t)}\right)^{n-2}\right) = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right),$$

$$\begin{aligned} \frac{1}{y'(t)^{n-2}}\int_t^{S_3}\rho_3V_1(s)\mathrm{d}s &= \frac{1}{y'(t)^{n-2}}\left[(y'(s)^{n-2} - z'(s)^{n-2})V_1V_2'\right]_t^{S_3} \\ &\quad -\frac{1}{y'(t)^{n-2}}\int_t^{S_3}(y'(s)^{n-2} - z'(s)^{n-2})V_1'V_2'\mathrm{d}s \\ &= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}(1 + |V_1(t)|) + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\int_t^{S_3}V_2'(s)\mathrm{d}s\right)\right) \\ &= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}|V_1(t)|\right). \end{aligned} \tag{4.55}$$

From (4.51), we see that $V_2(t) \leq -C_3$ for $t \in [S_4, S_3]$. Also since $V_1(S_3) < 0$, we have

$$V_1(t) \leq V_1(S_3) = V_2(S_3) + \mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right) \leq \frac{-C_3}{2}.$$

This gives

$$0 < \frac{1}{V_1(t)V_2(t)} \leq \frac{4}{C_3^2} = \mathcal{O}(1), \quad \frac{1}{|V_1(t)|} \leq \frac{2}{C_3} = \mathcal{O}(1). \tag{4.56}$$

From Lemma 4.2, (4.17), (4.53), (4.54) and (4.56) we have

$$\begin{aligned} V_1'(t)V_2(t) - V_1(t)V_2'(t) &= \left(\frac{y'(S_3)}{y'(t)}\right)^{n-2} (V_1'(S_3)V_2(S_3) - V_1(S_3)V_2'(S_3)) + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}|V_1(t)|\right) \\ &= \left(\frac{y'(S_3)}{y'(t)}\right)^{n-2} ((V_1'(S_3) - V_2'(S_3))V_1(S_3) + (V_2(S_3) - V_1(S_3))V_2'(S_3)) + \\ &\quad + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}|V_1(t)|\right) \\ &= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}|V_1(t)|\right). \end{aligned}$$

Dividing by V_1V_2 and integrating between $[t, S_3]$, we obtain

$$\frac{V_1'}{V_1} - \frac{V_2'}{V_2} = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\frac{1}{|V_2(t)|}\right) = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right),$$

$$V_1(t) = V_2(t)\frac{V_1(S_3)}{V_2(S_3)}e^{\mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)}$$

$$V_1(t) - V_2(t) = -\frac{V_2(t)}{V_2(S_3)}(V_2(S_3) - V_1(S_3)) + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^3}\right) = \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right).$$

From (4.29) and (4.30) we have

$$|V_1'(t) - V_2'(t)| = \mathcal{O}\left(|V_1'(S_3) - V_2'(S_3)| + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)\right) = \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right).$$

This proves the lemma. ☐

As an immediate consequence of this, we have the following

Corollary 4.4. *There exists a constant $C_4 \geq 0$ such that for γ large*

$$S \leq T_1 + (n-1)\log\left(\frac{g''}{(g')^2}\right) + 3(n-1)\log(\log(g')) + (n-1)C_4. \tag{4.57}$$

Proof. Let $S_5 := T_1 + (n-1)\log\left(\frac{g''}{(g')^2}\right) + 3(n-1)\log(\log(g'))$ and $X(t) \overset{\text{def}}{=} e^{\frac{T_1 - t}{n-1}}$. Then,

$$\begin{aligned} V_2'(S_5) &= \frac{n}{(n-1)^2}\frac{1}{X(S_5)} + \mathcal{O}\left(\frac{1}{X(S_5)^2}\right) \\ &= \frac{n}{(n-1)^2}\frac{g''}{(g')^2}(\log(g'))^3 + \mathcal{O}\left(\left(\frac{g''}{(g')^2}\right)^2(\log(g'))^6\right). \end{aligned}$$

If $S \leq S_5$, then we can trivially take $C_4 = 0$.

If $S > S_5$, then from Lemma 4.3, there exists a $C_5 > 0$ such that

$$V_2'(S) = V_2'(S) - V_1'(S) \leq C_5 \frac{(\log(g'))^3 g''}{(g')^2}. \tag{4.58}$$

Since $V_2'(S) = \mathcal{O}\left(\frac{g''\delta^3}{(g')^2}\right) \to 0$ as $\gamma \to \infty$, we must have

$$V_2'(S) = \frac{n}{(n-1)^2} \frac{X(S)}{(1+X(S))^2} \to 0 \text{ as } \gamma \to \infty. \tag{4.59}$$

Claim 4. We must have $S < T_1$ and hence $X(S) \geq 1$.

Proof of Claim 4. Suppose $S \in [S_3, S_0]$, then we must have $X(S_0) \leq X(S) \leq X(S_3)$. From the definition of S_0 and Lemma 4.2, we see that

$$(n-1) \leq X(S) \leq e^{k_1}$$

which is a contradiction to (4.59). Thus $S < S_3$ and hence trivially $S < T_1$ which proves the claim. □

Using (4.58) along with (4.59), we see that

$$V_2'(S) \geq C \frac{1}{X(S)} \text{ for } \gamma \text{ large.}$$

This implies that there is a constant $C_4 > 0$ such that

$$S \leq T_1 + (n-1)\log\left(\frac{g''}{(g')^2}\right) + 3(n-1)\log\log g' + C_4.$$

□

The estimates so far obtained are rough estimates and we need to improve them in order to prove the theorem. For this, we need some explicit formulas as follows:

Lemma 4.5. *Let* $X(t) = e^{\frac{T_1 - t}{n-1}}$, *then*

$$\begin{aligned} I_1(t) &= \int_t^\infty (z')^{n-2} z'' V_2' \mathrm{d}s \\ &= \frac{g'}{(n-1)} \left(\frac{n}{(n-1)g'}\right)^n \left[-\frac{1}{n(n-1)} + \sum_{r=0}^{n-1} \frac{(-1)^r C_r^{n-1}}{(r+2)(1+X(t))^{r+2}}\right], \end{aligned}$$

$$I_2(t) = \int_t^\infty (z')^n V_2' \mathrm{d}s = \frac{g'}{(n+1)} \left(\frac{n}{(n-1)g'}\right)^n \frac{X(t)^{n+1}}{(1+X(t))^{n+1}}$$

and

$$\begin{aligned} I(t) &= \frac{g''}{g'} I_1(t) + g'' I_2(t) \\ &= \left(\frac{n}{(n-1)g'}\right)^n g'' \left[\frac{1}{n} + \frac{1}{n-1} \sum_{r=0}^{n-1} \frac{(-1)^r C_r^{n-1}}{(r+2)(1+X(t))^{r+2}}\right] \\ &+ \left(\frac{n}{(n-1)g'}\right)^n g'' \left[\frac{n}{(n^2-1)} \frac{X(t)^{n+1} - (1+X(t))^{n+1}}{(1+X(t))^{n+1}}\right]. \end{aligned} \tag{4.60}$$

Proof. Since $\mathrm{d}X = -\dfrac{X}{n-1}\mathrm{d}t$ and $V_2' = -g'z''$. Then from (3.29)

$$\begin{aligned}
I_1 &= \int_t^\infty (z')^{n-2}z''V_2'\mathrm{d}s = -g'\int_t^\infty (z')^{n-2}(z'')^2\mathrm{d}s \\
&= -\frac{g'}{(n-1)^2}\left(\frac{n}{(n-1)g'}\right)^n \int_t^\infty \frac{X(s)^n}{(1+X(s))^{n+2}}\mathrm{d}s \\
&= -\frac{g'}{(n-1)}\left(\frac{n}{(n-1)g'}\right)^n \int_0^{X(t)} \frac{X^{n-1}}{(1+X)^{n+2}}\mathrm{d}X \\
&= -\frac{g'}{(n-1)}\left(\frac{n}{(n-1)g'}\right)^n \sum_{r=0}^{n-1}(-1)^r C_r^{n-1}\int_0^{X(t)} \frac{\mathrm{d}X}{(1+X)^{r+3}} \\
&= \frac{g'}{(n-1)}\left(\frac{n}{(n-1)g'}\right)^n \left[-\frac{1}{n(n+1)} + \sum_{r=0}^{n-1}\frac{(-1)^r C_r^{n-1}}{(r+2)(1+X(t))^{r+2}}\right].
\end{aligned}$$

$$\begin{aligned}
I_2(t) &= \int_t^\infty (z')^n V_2'\mathrm{d}s = -g'\int_t^\infty (z')^n z''\mathrm{d}s \\
&= \frac{g'}{n+1}z'(t)^{n+1} = \frac{g'}{n+1}\left(\frac{n}{(n-1)g'}\right)^{n+1}\frac{X(t)^{n+1}}{(1+X(t))^{n+1}}.
\end{aligned}$$

$$\begin{aligned}
I(t) &= \frac{g''}{g'}I_1(t) + g''I_2(t) \\
&= \left(\frac{n}{(n-1)g'}\right)^n g''\left[-\frac{1}{n(n^2-1)} + \frac{1}{(n-1)}\sum_{r=0}^{n-1}\frac{(-1)^r C_r^{n-1}}{(r+2)(1+X(t))^{n+2}}\right] \\
&+ g''\left(\frac{n}{(n-1)g'}\right)^n\left[\frac{n}{(n^2-1)}\frac{X(t)^{n+1}}{(1+X(t))^{n+1}}\right].
\end{aligned}$$

Since $-\dfrac{1}{n(n^2-1)} + \dfrac{n}{(n^2-1)} = \dfrac{1}{n}$, the lemma follows. ☐

Lemma 4.6. *Let*

$$J(t) = \left(1 - g'(y(t))y'(t) - \frac{g''(y(t))}{g'(y(t))}\right)y'(t)^{n-2}V_1'(t) + ((y')^{n-2}V_1')'.$$

Then for $a < b$

$$\begin{aligned}
J(a) &= J(b) + \int_a^b \frac{g''(y(s))}{g'(y(s))}y''(s)y'(s)^{n-2}V_1'(s)\mathrm{d}s \\
&+ \int_a^b \left(g''(y(s)) + \frac{g'''(y(s))}{g'(y(s))} - \left(\frac{g''(y(s))}{g'(y(s))}\right)^2\right)y'(s)^n V_1'(s)\mathrm{d}s. \qquad (4.61)
\end{aligned}$$

Proof. Multiply the equation (4.10) by $(g'(y)y' + \dfrac{g''(y)y'}{g'(y)} - 1)$ and integrate to obtain

$$\begin{aligned}
-\int_a^b (g'(y)y' + \frac{g''(y)}{g'(y)}y' - 1)((y')^{n-2}V_1')'\mathrm{d}s &= \frac{1}{n-1}\int_a^b (e^{g(y)+\log(g'(y))-s})'V_1\mathrm{d}s \\
&= \left[\frac{V_1}{n-1}g'(y)e^{g(y)-s}\right]_a^b - \frac{1}{n-1}\int_a^b g'(y)e^{g(y)-s}V_1'\mathrm{d}s
\end{aligned}$$

and

$$\begin{aligned}-\int_a^b (g'(y)y' + \frac{g''(y)}{g'(y)}y' - 1)((y')^{n-2}V_1')'\mathrm{d}s &= -\left[(g'(y)y' + \frac{g''(y)}{g'(y)}y' - 1)(y')^{n-2}V_1'\right]_a^b \\ &\quad + \int_a^b (g''(y) + \frac{g'''(y)}{g'(y)} - \left(\frac{g''(y)}{g'(y)}\right)^2)(y')^n V_1'\mathrm{d}s \\ &\quad + \int_a^b (g'(y) + \frac{g''(y)}{g'(y)})y''(y')^{n-2}V_1'\mathrm{d}s.\end{aligned}$$

Since $(n-1)g'(y)y''(y')^{n-2} = ((y')^{n-1})'g'(y) = -g'(y)e^{g(y)-s}$ hence cancelling this term on both sides yields the identity. ☐

Lemma 4.7. *Let $\delta > 0$, $\delta = \mathcal{O}(\log(g'))$ and $T_\delta = T_1 - (n-1)\delta$. For $t \geq \max\{S, T_\delta\} \overset{\text{def}}{=} S_4$ and $X(t) = e^{\frac{T_1 - t}{n-1}}$,*

$$\begin{aligned}L(t) &\overset{\text{def}}{=} \int_t^\infty \frac{g''(y)}{g'(y)}y''(y')^{n-2}V_1'\mathrm{d}s + \int_t^\infty (g''(y) + \frac{g'''(y)}{g'(y)} - \left(\frac{g''(y)}{g'(y)}\right)^2)(y')^n V_1'\mathrm{d}s \\ &= g''\left(\frac{n}{(n-1)g'}\right)^n \left[\frac{1}{n} + \frac{1}{(n-1)}\sum_{r=0}^{n-1}\frac{(-1)^r C_r^{n-1}}{(r+2)(1+X(t))^{r+2}}\right] \\ &\quad + g''\left(\frac{n}{(n-1)g'}\right)^n \left[\frac{n}{(n^2-1)}\frac{X(t)^{n+1} - (1+X(t))^{n+1}}{(1+X(t))^{n+1}} + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right], \qquad (4.62)\end{aligned}$$

then $J(t) = L(t)$. Furthermore, if $t \leq S_3 = S_1 - (n-1)k_1$, then

$$((y')^{n-2}V_1')'(t) = e^{\mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)}((z')^{n-2}V_2')'(t). \qquad (4.63)$$

Proof. Let $S_4 \leq t \leq S_3$, then $V_1(t) \leq -\tilde{C}_3$ and hence from (4.18), we have

$$-y'' = \frac{e^{g(y)-t}}{(n-1)(y')^{n-2}} = e^{\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right)}\frac{e^{g+g'(z-\gamma)-t}}{(n-1)(z')^{n-2}} = -e^{\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right)}z''$$

and from (4.53) and (4.23), we have

$$-((y')^{n-2}V_1')' = \rho_1 e^{-t}V_1 = e^{\mathcal{O}\left(\frac{g''\delta^2}{(g')^2}\right)}\rho_2 e^{-t}(V_2 + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)) = -e^{\mathcal{O}\left(\frac{g''\delta^3}{(g')^2}\right)}((z')^{n-2}V_2')'.$$

which proves (4.63). From (4.18) to (4.23) and from Lemma 4.3, we have

$$\begin{aligned}L(t) &= e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}\left[\frac{g''}{g'}\int_t^\infty (z')^{n-2}z''(V_2' + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right))\mathrm{d}s\right. \\ &\quad \left. + g''\int_t^\infty (z')^n(V_2' + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right))\mathrm{d}s + \mathcal{O}\left(\frac{g'''}{g'} + \left(\frac{g''}{g'}\right)^2\right)\int_t^\infty (z')^n V_1'\mathrm{d}s\right] \\ &= e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}\left[\frac{g''}{g'}I_1(t) + g''I_2(t)\right] \\ &\quad + \mathcal{O}\left(\frac{\delta^3 (g'')^2}{(g')^3}\int_t^\infty (z')^{n-2}z''\mathrm{d}s + \frac{\delta^3 (g'')^2}{(g')^2}\int_t^\infty (z')^n\mathrm{d}s + \left(\frac{g''}{(g')^n} + \left(\frac{g''}{g'}\right)^2\right)\int_t^\infty (z')^n V_1'\mathrm{d}s\right).\end{aligned}$$

Furthermore,

$$\int_t^\infty (z')^{n-2}z''\mathrm{d}s = \frac{z'(t)^{n-1}}{n-1} = \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n-1}\right),$$

and

$$\begin{aligned}\int_t^\infty (z')^n\mathrm{d}s &= \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n-1}\int_t^\infty z'(s)\mathrm{d}s\right) = \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n-1}(\gamma - z(t))\right)\\ &= \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n}\log(1+X(t))\right) = \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n}\delta\right).\end{aligned}$$

From Lemma 4.3, for $t \geq \max\{S, T_\delta\}$, we have $V_1(t) = V_2(t) + \mathcal{O}\left(\frac{g''\delta^3}{(g')^2}\right) = \mathcal{O}(1)$. Therefore

$$\int_t^\infty (z')^n V_1'\mathrm{d}s = \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n}(1-V_1(t))\right) = \mathcal{O}\left(\left(\frac{n}{(n-1)g'}\right)^{n}\right).$$

Hence we get

$$L(t) = e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}I(t) + \left(\frac{n}{(n-1)g'}\right)^{n}\mathcal{O}\left(\frac{\delta^3(g'')^2}{(g')^2} + \frac{\delta^4(g'')^2}{(g')^2}\right).$$

Now (4.51) follows from (4.60) and this completes the proof of the lemma. ☐

We now introduce an assumption on S:

Assumption 4.8. Let

$$S_6 = S_6(\gamma) \overset{\text{def}}{=} T_1 - (\frac{4q}{q-1}+1)(n-1)\log(g'), \tag{4.64}$$

and now assume that there exists a sequence $\gamma_l \to \infty$ such that $S = S(\gamma_l)$ satisfies

$$S_6(\gamma_l) \leq S(\gamma_l) = S. \tag{4.65}$$

For the next part, we assume that S satisfies Assumption 4.8 and derive the asymptotics as $\gamma_l \to \infty$. By a slight abuse of notation, for the subsequent sections, we denote γ_l by γ, $S_6(\gamma_l)$ to be S_6 and $S(\gamma_l)$ by S. We suppress writing the subsequence l and mean γ is large to denote γ_l is large. Then we have the following crucial result.

Lemma 4.9. *Assume that S satisfies Assumption 4.8, then for γ large, we have*

$$S = T_1 + (n-1)\log\left(\frac{(n-1)g''}{(g')^2} + \mathcal{O}\left(\frac{\delta^4(g'')^2}{(g')^4}\right)\right) \tag{4.66}$$

and

$$V_2'(S) = \frac{n}{n-1}\frac{g''}{(g')^2}\left(1+\mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right). \tag{4.67}$$

Proof. As before, we let $X(t) := e^{\frac{T_1-t}{n-1}}$. Then from (4.57), we trivially have

$$X(S) \geq \frac{e^{-C_4}(g')^2}{(\log(g'))^3 g''} \to \infty \quad \text{as } \gamma \to \infty.$$

We have

$$(z')^{n-2}V_2' = \frac{n}{(n-1)^2}\left(\frac{n}{(n-1)g'}\right)^{n-2}\frac{X(t)^{n-1}}{(1+X(t))^n},$$

$$\begin{aligned}((z')^{n-2}V_2')' &= \frac{n}{(n-1)^3}\left(\frac{n}{(n-1)g'}\right)^{n-2}\left(\frac{nX(t)^n}{(1+X(t))^{n+1}}-\frac{(n-1)X(t)^{n-1}}{(1+X(t))^n}\right)\\ &= \frac{n}{(n-1)^3}\left(\frac{n}{(n-1)g'}\right)^{n-2}\left(\frac{1}{X(t)}+\mathcal{O}\left(\frac{1}{(X(t))^2}\right)\right). \qquad (4.68)\end{aligned}$$

Set $a = S$ and $b = \infty$ in (4.61) and from (4.62), (4.63), (4.68) we have for $\delta = \log(g')$

$$\begin{aligned}((z')^{n-2}V_2')'(S) &= \frac{n}{(n-1)^3}\left(\frac{n}{(n-1)g'}\right)^{n-2}\left(\frac{1}{X(S)}+\mathcal{O}\left(\frac{1}{X(S)^2}\right)\right)\\ &= e^{\mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)}((y')^{n-2}V_1')'(S) = e^{\mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)}L(S)\\ &= e^{\mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)}g''\left((\frac{n}{(n-1)g'}\right)^{n}\left(\frac{1}{n}+\mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right)\end{aligned}$$

and

$$\frac{1}{X(S)} = \frac{(n-1)g''}{(g')^2}(1+\mathcal{O}\left(\frac{1}{X(S)}\right))+\mathcal{O}\left(\frac{1}{X(S)^2}+\delta^4\left(\frac{g''}{(g')^2}\right)^2\right).$$

Hence

$$e^{\frac{S-T_1}{n-1}} = \frac{1}{X(S)} = \frac{(n-1)g''}{(g')^2}\left[1+\mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right],$$

which gives after taking logarithms

$$S = T_1 + (n-1)\log\left(\frac{(n-1)g''}{(g')^2}+\mathcal{O}\left(\delta^4\left(\frac{g''}{(g')^2}\right)^2\right)\right)$$

and

$$\begin{aligned}V_2'(S) &= \frac{n}{(n-1)^2}\frac{X(S)}{(1+X(S))^2} = \frac{n}{(n-1)^2}\left(\frac{1}{X(S)}-\frac{2}{X(S)^2}+\mathcal{O}\left(\frac{1}{X(S)^3}\right)\right)\\ &= \frac{n}{n-1}\frac{g''}{(g')^2}\left(1+\mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right).\end{aligned}$$

This proves the lemma. ☐

Next we study the behaviour of V_1 for $t \in [T(\gamma), S]$.

Lemma 4.10. *Let S satisfies the Assumption 4.8, then there exists a $C_0 > 0$ such that for γ large with $t \in [S - C_0 g, S)$, the following holds:*

$$V_1(t) < 0, \quad \textit{and} \quad V_1'(t) < 0. \qquad (4.69)$$

Proof. Let S_6 defined in (4.64) and

$$t_0 \overset{\text{def}}{=} \inf\{t \in [T(\gamma), S];\ V_1(s) < 0 \text{ for } s \in (t, S]\}. \qquad (4.70)$$

Observe that from (4.66), there exists a $C_1 > 0$ such that for γ large $\tilde{T} \le S - C_1 g$.

Case i: If $\tilde{T} \ge t_0$, then there is nothing to prove.

Case ii: Assume that $\tilde{T} < t_0$. Then $V_1(t_0) = 0$ and $0 \le -V_1(t) \le -V_1(S)$ for $t \in (t_0, S)$. Hence from (3.13), (3.37), (4.9) and (4.10), we have

$$\begin{aligned}\int_{t_0}^{S_6}\rho_1 e^{-s}V_1\left(\int_{t_0}^{s}\frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right)\mathrm{d}s &= \mathcal{O}\left(|V_1(S)|(g')^{n-1}(S-t_0)\int_{\tilde{T}}^{S_6}e^{g(y(s))-s}\mathrm{d}s\right)\\ &= \mathcal{O}\left(\frac{|V_1(S)|(S-t_0)(g')^{n-1+\frac{q}{q-1}}}{(g')^{\frac{4q}{q-1}+n}}\right) = \mathcal{O}\left(\frac{|V_1(S)|(S-t_0)}{g}\right).\end{aligned}$$

Similarly, now from (4.18), we have

$$\begin{aligned}
\int_{S_6}^{S} \rho_1 e^{-s} V_1 \left(\int_{t_0}^{s} \frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right) \mathrm{d}s &= \mathcal{O}\left(|V_1(S)|(S-T_0)(g')^{n-1}\int_{S_6}^{S} e^{g+g'(z-\gamma)-s}\mathrm{d}s\right)\\
&= \mathcal{O}\left(|V_1(S)|(S-t_0)(g')^{n-1}|z'(S)^{n-1}-z'(S_6)^{n-1}|\right)\\
&= \mathcal{O}\left(|V_1(S)|(S-t_0)\left|\frac{X(S)^{n-1}}{(1+X(S))^{n-1}}-\frac{X(S_6)^{n-1}}{(1+X(S_6))^{n-1}}\right|\right)\\
&= \mathcal{O}\left(\frac{|V_1(S)|(S-t_0)}{X(S)}\right) = \mathcal{O}\left(\frac{|V_1(S)|(S-t_0)g''}{(g')^2}\right)\\
&= \mathcal{O}\left(\frac{|V_1(S)|(S-t_0)}{g}\right).
\end{aligned}$$

Here we have used the fact that for any $a>0$ and $b>0$, we have

$$\begin{aligned}
\frac{a^{n-1}}{(1+a)^{n-1}} - \frac{b^{n-1}}{(1+b)^{n-1}} &= \mathcal{O}\left(\frac{a^{n-1}(1+b)^{n-1}-b^{n-1}(1+a)^{n-1}}{(1+a)^{n-1}(1+b)^{n-1}}\right)\\
&= \mathcal{O}\left(\frac{a^{n-1}b^{n-2}+b^{n-1}a^{n-2}}{a^{n-1}b^{n-1}}\right) = \mathcal{O}\left(\frac{1}{a}+\frac{1}{b}\right) = \mathcal{O}\left(\frac{1}{a}\right).
\end{aligned}$$

Combining all the above estimates, we get

$$\begin{aligned}
|V_1(S)| &= -V_1(S)+V_1(t_0)\\
&= -\int_{t_0}^{S_6} \rho_1 V_1 e^{-s}\left(\int_{t_0}^{s}\frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right)\mathrm{d}s - \int_{S_6}^{S}\rho_1 V_1 e^{-s}\left(\int_{t_0}^{s}\frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right)\mathrm{d}s\\
&= \mathcal{O}\left(\frac{|V_1(S)|(S-t_0)}{g}\right).
\end{aligned}$$

This completes the proof of the lemma. ☐

Next we have

Lemma 4.11. *Let S satisfies the Assumption 4.8 and S_6 be as in* (4.64)*, then the following holds:*

$$V_1(S_6) = V_1(S)\left(1+(n-1)^2V_2'(S)-(n-1)(S-S_6)V_2'(S)\right)+\mathcal{O}\left(\frac{\delta^4 g''}{(g')^4}\right), \tag{4.71}$$

$$V_1'(S_6) = (n-1)V_1(S)V_2'(S)+\mathcal{O}\left(\frac{\delta^3 g''}{(g')^4}\right). \tag{4.72}$$

Proof. Let $t\in[S_6,S]$, then $X(t)=e^{\frac{T_1-t}{n-1}}\to\infty$ as $\gamma\to\infty$. From Lemma 4.10, $V_1(t)<0$ for $t\in[S_6,S]$, hence

$$\begin{aligned}
\log(-(n-1)V_2(t)) &= \log\left(1-\frac{n}{1+X(t)}\right) = -\frac{n}{1+X(t)}+\mathcal{O}\left(\frac{1}{X(t)^2}\right)\\
&= -(n-1)^2\frac{n}{(n-1)^2}\frac{X(t)}{(1+X(t))^2}+\mathcal{O}\left(\frac{1}{X(t)^2}\right)\\
&= -(n-1)^2V_2'(t)+\mathcal{O}\left(\frac{1}{X(t)^2}\right),
\end{aligned}$$

and $V_2'(S_6) = \mathcal{O}\left(\frac{1}{(g')^{\frac{4q}{q-1}+1}}\right)$. Therefore, from lemma 4.3,

$$\begin{aligned}
\int_{S_6}^{S} \frac{V_1(s)V_2'(s)}{V_2(s)}\mathrm{d}s &= \int_{S_6}^{S}(\log(-(n-1)V_2(s))'V_1(s)\mathrm{d}s \\
&= [V_1(s)\log(-(n-1)V_2(s)]_{S_6}^{S} - \int_{S_6}^{S} V_1'(s)\log(-(n-1)V_2(s))\mathrm{d}s \\
&= -(n-1)^2V_1(S)V_2'(S) + \mathcal{O}\left(V_1(S_6)V_2'(S_6)\right) + \mathcal{O}\left(V_1'(S_6)\int_{S_6}^{S} V_2'(s)\mathrm{d}s\right) \\
&= -(n-1)^2V_1(S)V_2'(S) + \mathcal{O}\left(\frac{V_1(S_6)}{(g')^{\frac{4q}{q-1}+1}}\right) + \mathcal{O}\left(V_1'(S_6)(V_2(S)-V_2(S_6))\right).
\end{aligned}$$

Since $V_2(S) - V_2(S_6) = \frac{n}{n-1}\left(\frac{1}{1+X(S)} - \frac{1}{1+X(S_6)}\right) = \mathcal{O}\left(\frac{1}{X(S)}\right)$, hence from lemma 4.9,

$$\int_{S_6}^{S} \frac{V_1(s)V_2'(s)}{V_2(s)}\mathrm{d}s = -(n-1)^2V_1(S)V_2'(S) + \mathcal{O}\left(\frac{V_1'(S_6)g''}{(g')^2} + \frac{(V_1(S_6))}{(g')^{\frac{4q}{q-1}+1}}\right). \tag{4.73}$$

With $\delta = \log(g')$ for $t \in [S_6, S]$, we have

$$y'(t) = e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}z'(t), \quad y''(t) = -\frac{e^{g(y)-t}}{(n-1)(y')^{n-2}} = e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}z''(t),$$

$$\begin{aligned}
y'(t)^{n-3}y''(t) - z'(t)^{n-3}z''(t) &= \left(1 - e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}\right)(z')^{n-3}z'' \\
&= \mathcal{O}\left(\frac{\delta^2 g''}{(g')^n}\frac{1}{X(S)}\right) = \mathcal{O}\left(\frac{\delta^2 (g'')^2}{(g')^{n+2}}\right),
\end{aligned}$$

$$y'(t)^{n-2} - z'(t)^{n-2} = \left(1 - e^{\mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\right)}\right)z'(t)^{n-2} = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^n}\right)$$

and

$$V_2'(t) + |V_2''(t)| = \mathcal{O}\left(\frac{1}{X(S)}\right) = \mathcal{O}\left(\frac{g''}{(g')^2}\right).$$

Hence

$$\begin{aligned}
\rho_3 &= ((y'(t)^{n-2} - z'(t)^{n-2})V_2')' \\
&= (y'(t)^{n-2} - z'(t)^{n-2})V_2'' + (n-2)(y'(t)^{n-3}y''(t) - z'(t)^{n-3}z''(t)V_2' = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^{n+2}}\right).
\end{aligned}$$

and

$$\frac{1}{y'(t)^{n-2}}\int_t^S \rho_3 V_1 \mathrm{d}s = \mathcal{O}\left(\frac{(S-S_6)\delta^2 g''}{(g')^4}\right) = \mathcal{O}\left(\frac{\delta^3 g''}{(g')^4}\right).$$

From (4.23),

$$\frac{1}{y'(t)^{n-2}}\int_t^S (\rho_1 - \rho_2)V_1V_2e^{-s}\mathrm{d}s = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}\frac{1}{y'(t)^{n-2}}\int_t^S \rho_1 V_1 e^{-s}\mathrm{d}s\right) = \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}V_1'(t)\right).$$

Since $V_2'(S_6) = \mathcal{O}\left(\frac{1}{X(S_6)}\right) = \mathcal{O}\left(\frac{1}{(g')^{\frac{4q}{q-1}+1}}\right)$, we have from (4.17),

$$\begin{aligned}
V_1'(t) &= \frac{V_1(t)V_2'(t)}{V_2(t)} - \left(\frac{y'(S)}{y'(t)}\right)^{n-2} \frac{V_1(S)V_2'(S)}{V_2(t)} + \frac{1}{y'(t)^{n-2}V_2(t)} \int_t^S (\rho_1 - \rho_2)V_1V_2e^{-s}\mathrm{d}s \\
&\quad + \frac{1}{y'(t)^{n-2}V_2(t)} \int_t^S \rho_3 V_1 \mathrm{d}s \\
&= \frac{V_1(t)V_2'(t)}{V_2(t)} + (n-1)V_1(S)V_2'(S) + \mathcal{O}\left(\frac{\delta^2 g''}{(g')^2}V_1'(t) + \frac{\delta^3 g''}{(g')^4}\right).
\end{aligned}$$

Hence solving for $V_1'(t)$ to obtain

$$V_1'(t) = \frac{V_1(t)V_2'(t)}{V_2(t)} + (n-1)V_1(S)V_2'(S) + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^4}\right). \tag{4.74}$$

Thus we get

$$V_1'(S_6) = (n-1)V_1(S)V_2'(S) + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^4}\right)$$

and this proves (4.72). Integrating (4.74) and using the above estimates to obtain

$$\begin{aligned}
V_1(S) - V_1(S_6) &= \int_{S_6}^S V_1(t)\frac{V_2'(t)}{V_2(t)}\mathrm{d}t + (n-1)V_1(S)V_2'(S)(S - S_6) + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^4}\right) \\
&= -(n-1)^2 V_1(S)V_2'(S) + (n-1)V_1(S)V_2'(S)(S-S_6) + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^4}\right).
\end{aligned}$$

This proves (4.71) and hence the lemma. ☐Next, we have

Lemma 4.12. *Assume S satisfies* (4.65). *Then for $t \in [\tilde{T}, S_6]$*

$$V_1'(t) = V_1'(S_6) + \mathcal{O}\left(\frac{(g'')^2}{(g')^4}\right), \tag{4.75}$$

$$V_1(\tilde{T}) = V_1(S_6) + V_1'(S_6)(\tilde{T} - S_6) + \mathcal{O}\left(\frac{g''}{(g')^2}\right). \tag{4.76}$$

Proof. From Lemma 4.10, $V_1(t) < 0$, $V_1'(t) < 0$ for $t \geq S - C_0 g$. Let t_0 as in (4.70) and define $t_1 \leq t_0 < S_6$ by

$$t_1 \overset{\text{def}}{=} \inf\{t < t_0;\ V_1(s) > 0, V_1'(s) < 0 \text{ for } s \in (t, t_0)\}, \tag{4.77}$$

that is t_0 is the first zero before S and t_1 is the second turning point of V_1 if it exists.

Claim 5. For $t \in [\tilde{T}, S_6]$, we have $V_1(t) = \mathcal{O}(1)$ and $V_1'(t) \leq 0$.

Proof of Claim 5. Assume that $t_1 \geq \tilde{T}$ exists. Then either (i) $V_1'(t_1) = 0$ or (ii) $t_1 = \tilde{T}$ and $V_1(\tilde{T}) > 0$, $V_1'(t) < 0$ for $t \in [\tilde{T}, S_6]$. From (4.72), we have $V_1'(S_6) = \mathcal{O}\left(\frac{g''}{(g')^2}\right)$ and

$$V_1'(S_6)\int_t^{S_6} \left(\frac{y'(S_6)}{y'(s)}\right)^{n-2} \mathrm{d}s = \mathcal{O}\left(\frac{g''(S_6 - t)}{(g')^2}\right) = \mathcal{O}\left(\frac{g''(S_6 - \tilde{T})}{(g')^2}\right) = \mathcal{O}\left(\frac{g''g}{(g')^2}\right) = \mathcal{O}(1).$$

From (4.10) and (3.37), we have for $t_1 \leq a \leq b \leq S_6$

$$\begin{aligned}
&\int_a^b \rho_1 V_1 e^{-s}\left(\int_a^s \frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right)\mathrm{d}s = \mathcal{O}\left(\max\left\{|V_1(a)|,|V_1(b)|\right\}(g')^{n-1}\int_{\tilde{T}}^{S_6} e^{g(y(s))-s}\mathrm{d}s\right)\\
= \;& \max\left\{|V_1(a)|,|V_1(b)|\right\}\mathcal{O}\left(\frac{(\tilde{T}-S_6)^2(g')^{n-1}}{(g')^{\frac{2q}{q-1}+n}}\right) = \max\left\{|V_1(a)|,|V_1(b)|\right\}\mathcal{O}\left(\frac{1}{(g')^{\frac{q}{q-1}+1}}\right).
\end{aligned}$$

Hence for $t \in [t_1, S_6]$

$$\begin{aligned}
V_1(t) &= V_1(S_6) - V_1'(S_6)\int_t^{S_6}\left(\frac{y'(S_6)}{y'(\theta)}\right)^{n-2}\mathrm{d}\theta - \int_t^{S_6}\rho_1 V_1 e^{-s}\left(\int_t^s \frac{\mathrm{d}\theta}{y'(\theta)^{n-2}}\right)\mathrm{d}s\\
&= \mathcal{O}(1) + \max\left\{|V_1(t)|,|V_1(S_6)|\right\}\mathcal{O}\left(\frac{1}{(g')^{\frac{q}{q-1}+1}}\right).
\end{aligned}$$

This implies $V_1(t) = \mathcal{O}(1)$. Suppose $V_1'(t_1) = 0$, then taking $a = t_1$, $b = t_0$ to obtain

$$0 = V_1(t_0) = V_1(t_1) - \int_{t_1}^{t_0}\rho_1 V_1 e^{-s}\left(\int_t^s \frac{\partial\theta}{(y'(\theta))^{n-2}}\right)\mathrm{d}s = V_1(t_1) + \mathcal{O}\left(\frac{V_1(t_1)}{(g')^{\frac{q}{q-1}+1}}\right)$$

which gives a contradiction if $V_1(t_1) > 0$ and this proves the claim. ☐

From (3.37), (3.39), (4.12), (4.72) and the above claim, we have for $t \in [\tilde{T}, S_6]$

$$\begin{aligned}
V_1'(t) &= \left(\frac{y'(S_6)}{y'(t)}\right)^{n-2}V_1'(S_6) + \frac{1}{y'(t)^{n-2}}\int_t^{S_6}\rho_1 V_1 e^{-s}\mathrm{d}s\\
&= \left(\frac{1+\mathcal{O}\left(\frac{g''}{(g')^2}\right)}{1+\mathcal{O}\left(\frac{g''}{(g')^2}\right)}\right)V_1'(S_6) + \mathcal{O}\left((g')^{n-1}\int_t^{S_6} e^{g(y(s))-s}\mathrm{d}s\right) = V_1'(S_6) + \mathcal{O}\left(\frac{(g'')^2}{(g')^4}\right).
\end{aligned}$$

Integrating the above expression, we get

$$\begin{aligned}
V_1(\tilde{T}) &= V_1(S_6) + V_1'(S_6)(\tilde{T}-S_6) + \mathcal{O}\left(\frac{(\tilde{T}-S_6)(g'')^2}{(g')^4}\right)\\
&= V_1(S_6) + V_1'(S_6)(\tilde{T}-S_6) + \mathcal{O}\left(\frac{\gamma g'(g'')^2}{(g')^4}\right)\\
&= V_1(S_6) + V_1'(S_6)(\tilde{T}-S_6) + \mathcal{O}\left(\frac{g''}{(g')^2}\right).
\end{aligned}$$

This proves the lemma. ☐

Lemma 4.13. *Assume that S satisfies Assumption 4.8, then for $t \in [T(\gamma), \tilde{T}]$, we have*

$$V_1'(t) = V_1'(\tilde{T}) + \mathcal{O}\left(\frac{(g'')^2}{(g')^3} + \frac{e^{-(g-\frac{(n-1)\gamma g'}{n})}}{(g')^{\alpha}}\right), \tag{4.78}$$

$$V_1(T(\gamma)) = V_1(\tilde{T}) + V_1'(\tilde{T})(T(\gamma)-\tilde{T}) + \mathcal{O}\left(\frac{g''}{(g')^3} + \frac{e^{-(g-\frac{(n-1)\gamma g'}{n})}}{(g')^{\alpha}}\right). \tag{4.79}$$

Proof. First we show that for $t \in [T(\gamma), \tilde{T}]$

$$V_1(t) = O(1), V_1'(t) < 0. \tag{4.80}$$

For this, let t_0 and t_1 as in (4.70) and in (4.77).

Then from Claim 5, it follows that $t_1 < \tilde{T}$. Then either $t_1 > T(\gamma)$ and $V_1'(t_1) = 0$, $V_1(t_1) > 0$ or $t_1 = T(\gamma)$ and $V_1'(t) < 0$ for all $T(\gamma) < t < S$. For $t \in [T(\gamma), \tilde{T}]$, $y(t) \leq s_0$ and hence $\rho_1(t) = \dfrac{f'(y(t))}{n-1} = \mathcal{O}\left(y(t)^{-1+\alpha}\right)$. Therefore for $t \in (T(\gamma), \tilde{T})$, there exists a $\xi \in (T(\gamma), t)$ such that $y(t) = y'(\xi)(t - T(\gamma))$. Hence from (1.4) and (3.48) we have $\rho_1(t) = \dfrac{f'(y(t))}{n-1} = \mathcal{O}\left(\dfrac{(t - T(\gamma))^{-1+\alpha}}{(g')^{-1+\alpha}}\right)$. Hence from (2.10), (2.11), (3.38), (4.13), (4.72) and (4.75), we have

(4.81)

$$\begin{aligned}
V_1(t) &= V_1(\tilde{T}) - V_1'(\tilde{T}) \int_t^{\tilde{T}} \left(\frac{y'(\tilde{T})}{y'(\theta)}\right)^{n-2} \mathrm{d}\theta - \int_t^{\tilde{T}} \rho_1 V_1 e^{-s} \left(\int_t^s \frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right) \mathrm{d}s \\
&= V_1(\tilde{T}) - V_1'(\tilde{T}) \int_t^{\tilde{T}} \left(\frac{1 + \mathcal{O}\left(\frac{g''}{(g')^2}\right)}{1 + \mathcal{O}\left(\frac{g''}{(g')^2}\right)}\right) \mathrm{d}\theta + (\max_{s \in [t, \tilde{T}]} |V_1(s)|) \mathcal{O}\left((g')^{n-1-\alpha} \int_{T(\gamma)}^{\tilde{T}} (s - T(\gamma))^{\alpha} e^{-s} \mathrm{d}s\right) \\
&= V_1(\tilde{T}) - V_1'(\tilde{T})(\tilde{T} - t) + \mathcal{O}\left(\left(\frac{g''}{(g')^2}\right)^2\right)(\tilde{T} - T(\gamma)) \\
&\quad + (\max_{s \in [t, \tilde{T}]} |V_1(s)|) \mathcal{O}\left((g')^{n-1-\alpha} \int_{T(\gamma)}^{\tilde{T}} (1 + s - T(\gamma)) e^{-s} \mathrm{d}s\right) \\
&= V_1(\tilde{T}) - V_1'(\tilde{T})(\tilde{T} - t) + \mathcal{O}\left(\frac{(g'')^2}{(g')^3} + \max_{s \in [t, \tilde{T}]} |V_1(s)| (g')^{n-\alpha-1} e^{-T(\gamma)}\right).
\end{aligned}$$

Therefore, we have

$$V_1(t) = V_1(\tilde{T}) - V_1'(\tilde{T})(\tilde{T} - t) + \mathcal{O}\left(\frac{(g'')^2}{(g')^3} + \frac{e^{-\left(g - \left(\frac{n-1}{n}\right)\gamma g'\right)}}{(g')^{\alpha}} \max_{s \in [t, \tilde{T}]} |V_1(s)|\right).$$

Since from Claim 5, $V_1(\tilde{T}) = \mathcal{O}(1)$, hence

$$V_1(t) = \mathcal{O}(1) \text{ for all } t \in [T(\gamma), \tilde{T}]. \tag{4.82}$$

Suppose $V_1'(t_1) = 0$ and $V_1(t_1) > 0$, then $V_1(t_0) = 0$ and

$$\begin{aligned}
0 &= V_1(t_0) = V_1(t_1) + V_1'(t_1)(t_0 - t_1) - \int_{t_1}^{t_0} \rho_1 V_1 e^{-s} \left(\int_{t_1}^s \frac{\mathrm{d}\theta}{(y'(\theta))^{n-2}}\right) \mathrm{d}s \\
&= V_1(t_1) + \mathcal{O}\left(\frac{e^{-(g - \frac{(n-1)\gamma g'}{n})}}{(g')^{\alpha}}\right) V_1(t_1)
\end{aligned}$$

which is a contradiction if $V_1(t_1) > 0$ and this proves (4.80).

From (4.81) and (4.82), the estimate

$$V_1(T(\gamma)) = V_1(\tilde{T}) + V_1'(\tilde{T})(T(\gamma) - \tilde{T}) + \mathcal{O}\left(\frac{g''}{(g')^3} + \frac{e^{-(g - \frac{(n-1)\gamma g'}{n})}}{(g')^{\alpha}}\right)$$

follows and this proves the lemma. □

Lemma 4.14. *Assume S satisfies* (4.65) *and* $\delta = \mathcal{O}(\log(g'))$. *Then*

$$V_1(T(\gamma)) = V_1(S)\left(1 - \frac{(n-1)\gamma g''}{(g')^2} + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^2} + \frac{\delta^3}{(g')^2}\right)\right), \tag{4.83}$$

$$T'(\gamma) = \frac{\left(1 + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right)\right)}{n}\left(g' - (n-1)\gamma g'' + \mathcal{O}\left(\frac{\delta^4 g''}{g'} + \frac{\delta^3}{g'}\right)\right). \tag{4.84}$$

Proof. Denote $\eta = \eta(\gamma) \overset{\text{def}}{=} \dfrac{e^{-(g - \frac{(n-1)\gamma g'}{n})}}{(g')^\alpha}$. Then from Lemmas 4.9 to 4.13, we have

$$\begin{aligned}
V_1(T(\gamma)) &= V_1(\tilde{T}) + V_1'(\tilde{T})(T(\gamma) - \tilde{T}) + \mathcal{O}\left(\frac{g''}{(g')^2} + \eta\right) \\
&= V_1(S_6) + V_1'(S_6)(\tilde{T} - S_6) + (V_1'(S_6) + \mathcal{O}\left(\frac{(g'')^2}{(g')^4}\right))(T(\gamma) - \tilde{T}) + \mathcal{O}\left(\frac{g''}{(g')^2} + \eta\right) \\
&= V_1(S_6) + V_1'(S_6)(T(\gamma) - S_6) + \left(\frac{g''}{(g')^2} + \eta\right) \\
&= V_1(S)\left(1 + (n-1)^2 V_2'(S) + (n-1)V_2'(S)(S_6 - S)\right) \\
&\quad + (n-1)V_1(S)V_2'(S)(T(\gamma) - S_6) + \mathcal{O}\left(\frac{\delta^3 g''}{(g')^4}(T(\gamma) - S_6) + \eta\right) \\
&= V_1(S)\left(1 + (n-1)^2 V_2'(S) + (n-1)V_2'(S)(T(\gamma) - S)\right) + \mathcal{O}\left(\frac{\delta^3}{(g')^2} + \eta\right).
\end{aligned}$$

Now, (2.11) and (4.66)

$$\begin{aligned}
T(\gamma) - S &= g - \frac{(n-1)\gamma g'}{n} + (n-1)\log(\frac{(n-1)g'}{n}) + \frac{(n-1)\alpha_n \gamma g''}{g'} + A(\gamma) \\
&\quad - g - (n-1)\log(\frac{(n-1)g'}{n}) - (n-1)\log(\frac{(n-1)g''}{(g')^2}) + \mathcal{O}\left(\frac{\delta g''}{(g')^2} + \frac{\delta^2}{g'}\right) \\
&= -\frac{(n-1)\gamma g'}{n} - (n-1)\log\left(\frac{(n-1)g''}{(g')^2}\right) + \mathcal{O}(1)
\end{aligned}$$

and from (4.67)

$$\begin{aligned}
(n-1)V_2'(S)(T(\gamma) - S) &= \frac{ng''}{(g')^2}\left(1 + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right)(-\frac{(n-1)\gamma g'}{n} - (n-1)\log(\frac{(n-1)g''}{(g')^2}) + \mathcal{O}(1)) \\
&= -\frac{(n-1)\gamma g''}{g'} - \frac{n(n-1)g''}{(g')^2}\log\left(\frac{g''}{(g')^2}\right) + \mathcal{O}\left(\frac{\delta^4 \gamma (g'')^2}{(g')^3} + \frac{\delta^5 (g'')^2}{(g')^4}\right) \\
&= -\frac{(n-1)\gamma g''}{g'} + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right).
\end{aligned}$$

Therefore,

$$V_1(T(\gamma)) = V_1(S)\left(1 + \frac{n(n-1)g''}{(g')^2} - \frac{(n-1)\gamma g''}{g'}\right) + \mathcal{O}\left(\frac{\delta^4 g''}{(g')^2} + \frac{g''\delta^3}{(g')^2} + \eta\right).$$

This proves (4.83). From (H2) of Hypothesis 1, (2.12), (3.48) and (4.80) we have

$$\begin{aligned} T'(\gamma) &= -\frac{V_1(T(\gamma))}{y'(T(\gamma))} = -V_1(S)(1+\mathcal{O}\left(\frac{1}{(g')^2}\right))\frac{(n-1)g'}{n}\left(1-\frac{(n-1)\gamma g''}{g'}+\mathcal{O}\left(\frac{\delta^4 g''}{(g')^2}\right)\right) \\ &= \frac{(1+\mathcal{O}\left(\frac{\delta^3 g''}{(g')^2}\right))}{n}\left(g'-(n-1)\gamma g''+\mathcal{O}\left(\frac{\delta^4 g''}{g'}\right)\right). \end{aligned}$$

This proves the lemma. ☐

Proof of Theorem 2.3. Let $S_6(\gamma)$ be as in (4.64). Choose a $\gamma_0 > 0$ large such that for $\gamma \geq \gamma_0$, all the estimates up to Lemma 4.7 hold. Let $S = S(\gamma)$ be the first turning point as defined earlier. Let

$$I_1 \overset{\text{def}}{=} \{\gamma \geq \gamma_0;\ S(\gamma) < S_6(\gamma)\}, \tag{4.85}$$

$$I_2 \overset{\text{def}}{=} \{\gamma \geq \gamma_0;\ S(\gamma) \geq S_6(\gamma)\}. \tag{4.86}$$

Since $\gamma \to S(\gamma)$ is continuous, we have that I_2 is a closed set. Furthermore from Lemma 4.9, there exists a $\gamma_1 \geq \gamma_0$ such that if $\gamma \geq \gamma_1$, $\gamma \in I_2$, then

$$S(\gamma) = T_1 + (n-1)\log\left(\frac{g''}{(g')^2}\right) + \mathcal{O}(1). \tag{4.87}$$

Hence if $I_3 \overset{\text{def}}{=} \{\gamma > \gamma_1;\ S(\gamma) \geq S_6(\gamma)\}$ is non empty, then I_3 is both open and closed in (γ_1, ∞) which implies that $I_3 = (\gamma_1, \infty)$. Therefore we have either $I_3 = (\gamma_1, \infty)$ or $I_3 = \emptyset$.

If $I_3 = \emptyset$, then

$$I_4 \overset{\text{def}}{=} \{\gamma \geq \gamma_1;\ S(\gamma) < S_6(\gamma)\} = [\gamma_1, \infty).$$

Claim 6. $I_4 = \emptyset$.

Proof of Claim 6. Suppose $I_4 \neq \emptyset$, then $I_4 = [\gamma_1, \infty)$. Let $\gamma \geq \gamma_1$ and t_0 be as in (4.70). Then, for $t_0 \leq t \leq S$, we have $|V_1(t)| \leq |V_1(S)|$. Also $X(S_6) = e^{\frac{T_1 - S_6}{n-1}} = (g')^{\frac{4q}{q-1}+1}$ and hence for $t \in [t_0, S_6(\gamma)]$,

$$z'(t) = \mathcal{O}\left(\frac{1}{g'}\right), \quad z''(t) = \mathcal{O}\left(\frac{1}{X(S_6)g'}\right) = \mathcal{O}\left(\frac{1}{(g')^{\frac{2q}{q-1}+2}}\right)$$

$$z'(t)^{n-3}z''(t) = \mathcal{O}\left(\frac{1}{(g')^{\frac{2q}{q-1}+n-1}}\right), \quad |V_2'(t)| + |V_2''(t)| = \mathcal{O}\left(\frac{1}{X(S_6)}\right) = \mathcal{O}\left(\frac{1}{(g')^{\frac{2q}{q-1}+1}}\right).$$

Therefore from (3.12), (3.16) and (3.39), we have the following estimates for $t \in [\max\{t_0, \tilde{T}\}, S_6(\gamma)]$:

$$\begin{aligned} \frac{1}{y'(t)^{n-2}}\int_t^S \rho_1 V_1 V_2 e^{-s}\mathrm{d}s &= \mathcal{O}\left(|V_1(S)|(g')^{n-1}\int_t^S e^{g+g'(z-\gamma)-s}\mathrm{d}s\right) \\ &= \mathcal{O}\left(|V_1(S)|(g')^{n-1}[(z'(S))^{n-1} - (z'(t))^{n-1}]\right) \\ &= \mathcal{O}\left(|V_1(S)|\frac{(g')^{n-1}}{X(S_6)(g')^{n-1}}\right) = \mathcal{O}\left(\frac{|V_1(S)|}{(g')^{\frac{4q}{q-1}+1}}\right), \end{aligned}$$

$$\frac{1}{y'(t)^{n-2}}\int_t^S \rho_2 V_1 V_2 e^{-s}\mathrm{d}s = \mathcal{O}\left(\frac{|V_1(S)|(g')^{n-2}}{(g')^{n-1}X(S_6)}\right) = \mathcal{O}\left(\frac{|V_1(S)|}{(g')^{\frac{2q}{q-1}+2}}\right).$$

From (3.35), (3.36) and(3.37), we have

$$-(n-1)(y')^{n-2}y'' = -((y')^{n-1})' = e^{g(y(t))-t} \le e^{\psi(y(t))} = \mathcal{O}\left(\frac{1}{(g')^{\frac{2q}{q-1}+n}}\right)$$

and hence we get

$$\rho_3(t) = ((y')^{n-2} - (z'(t))^{n-2})V_2'' + (n-2)\left(\frac{(y')^{n-2}y''}{y'} - \frac{(z')^{n-2}z''}{z'}\right)V_2' = \mathcal{O}\left(\frac{1}{(g')^{\frac{2q}{q-1}+n-1}}\right).$$

From the above estimates, (4.17) and the fact that $S_6 - \tilde{T} = \mathcal{O}(g)$, we get

$$V_1'(t)V_2(t) = V_1(t)V_2'(t) - \left(\frac{y'(S)}{y'(t)}\right)^{n-2} V_2'(S)V_1(S) + \mathcal{O}\left(\frac{|V_1(S)|g}{(g')^{\frac{2q}{q-1}+1}}\right) = \mathcal{O}\left(\frac{|V_1(S)|g}{(g')^{\frac{2q}{q-1}+1}}\right).$$

Since $V_2(t) \le -C$ for some $C > 0$ independent of γ, we thus see that

$$V_1'(t) = \mathcal{O}\left(\frac{|V_1(S)|g}{(g')^{\frac{2q}{q-1}+1}}\right) = \mathcal{O}\left(\frac{|V_1(S)|}{(g')^{\frac{q}{q-1}+1}}\right). \tag{4.88}$$

Now from (4.53), we have

$$\begin{aligned} V_1(t) &= V_1(S) + \mathcal{O}\left(\frac{|V_1(S)|g(\tilde{T}-S)}{(g')^{\frac{2q}{q-1}+1}}\right) \le V_1(S_6) + \mathcal{O}\left(\frac{|V_1(S)|g^2}{(g')^{\frac{2q}{q-1}+1}}\right) \\ &= V_2(S_6) + \mathcal{O}\left(\frac{(log(g'))^3 g''}{(g')^2} + \frac{|V_1(S)|}{g'}\right) \le -\frac{1}{n-1} + \mathcal{O}\left(\frac{g''\delta^4}{(g')^2}\right) < 0. \end{aligned}$$

Hence $t_0 < \tilde{T}$ and $V_1(t) = V_1(S)(1 + \mathcal{O}\left(\frac{1}{(g')^{\frac{q}{q-1}+1}}\right))$. Now for $t \in [t_0, \tilde{T}]$ and from (4.12) and (4.88), we have

$$\begin{aligned} V_1'(t) &= \left(\frac{y'(\tilde{T})}{y'(t)}\right)^{n-2} V_1'(\tilde{T}) + \frac{1}{y'(t)^{n-2}} \int_t^{\tilde{T}} f'(y)V_1 e^{-s}\mathrm{d}s \\ &= \mathcal{O}\left(\frac{|V_1(S)|}{(g')^{\frac{q}{q-1}+1}} + |V_1(S)|(g')^{n-2} \int_t^{\tilde{T}} f'(y(s))e^{-s}\mathrm{d}s\right). \end{aligned}$$

As in (4.81), we therefore have

$$\begin{aligned} V_1(t) &= V_1(\tilde{T}) + |V_1(S)|\mathcal{O}\left(\frac{\tilde{T} - T(\gamma)}{(g')^{\frac{q}{q-1}+1}} + |V_1(S)|e^{-T(\gamma)}(g')^{n-2}\right) \\ &= V_1(S)\left(1 + \mathcal{O}\left(\frac{1}{(g')^{\frac{q}{q-1}}} + \frac{e^{-(g-\frac{(n-1)\gamma g'}{n})}}{(g')^{\alpha}}\right)\right). \end{aligned}$$

Since from (H2) of Hypothesis 1 which gives $g - \dfrac{(n-1)\gamma g'}{n} \ge b$ for some $b \in \mathbb{R}$ as $\gamma \to \infty$, we hence get

$$V_1(t) = V_1(S)(1 + o(1)) < 0.$$

Therefore $t_0 = T(\gamma)$ and

$$\begin{aligned} T'(\gamma) &= -\frac{V_1(T(\gamma))}{y'(T(\gamma))} = -\frac{(n-1)g'}{n}V_1(S)(1+o(1)) \geq \frac{(n-1)g'}{n}|V_1(S_6)|(1+o(1)) \\ &= \frac{g'}{n}(1+o(1)). \end{aligned}$$

Integrating the above inequality from γ_1 to γ and using (2.11) in Theorem 2.2, we obtain

$$g - \left(\frac{n-1}{n}\right)\gamma g' + \mathcal{O}(\log(g')) = T(\gamma) \geq T(\gamma_1) + \mathcal{O}(g(\gamma_1)) + \frac{g}{n}(1+o(1)).$$

Therefore we get

$$n - \frac{(n-1)\gamma g'}{g} \geq 1 + o(1).$$

Since from (H1) of Hypothesis 1, we see that

$$(n-1)\left[1 - \frac{\gamma g'}{g}\right] = (n-1)\left[1 - q + \mathcal{O}\left(\frac{\rho'\gamma}{g} + \frac{\rho}{g}\right)\right] \geq o(1)$$

which implies that $q \leq 1$ as $\gamma \to \infty$ which is a contradiction. This proves the claim. ☐

Therefore $I_3 = [\gamma_1, \infty)$ and for $\gamma \in I_3$ and from (4.84) the theorem follows. ☐

Acknowledgement: Adimurthi and J. Giacomoni were funded by IFCAM (Indo-French Centre for Applied Mathematics) under the project “Singular phenomena in reaction diffusion equations and in conservation laws” .

References

[1] Adimurthi, Existence of positive solutions of the semi linear Dirichlet problem with critical growth for the n-Laplacian, Ann. Sc. Norm. Sup. Pisa, XVII (1990), 393–413.

[2] Adimurthi, Uniqueness of positive solutions of a quasilinear Dirichlet problem with exponential nonlinearity, Proc. Roy. Soc. A 128, 5 (1998), 895–906.

[3] Adimurthi and S. K. Prashanth, Critical exponent problem in $\mathbb{R}^2$-borderline between existence and nonexistence of positive solutions for Dirichlet problem, Adv. Differential Equations, 5 (2000), 67–95.

[4] Adimurthi and K. Sandeep, Asingular Moser-Trudinger embedding and its applications, NODEA Nonlinear Differential Equations Appl., 13(5-6) (2007), 583-603.

[5] Adimurthi and S. L. Yadava, An elementary proof of the uniqueness of positive radial solutions of a quasilinear Dirichlet problem, Arch. Rational Mech. Anal., 127(3) (1994), 219–229.

[6] W. Allegretto and Yin Xi Huang, A Picone's identity for the p-Laplacian and applications, Nonlinear Anal., 32(7) (1998), 819-830.

[7] F. V. Atkinson and L. A. Peletier, Ground states and Dirichlet problems for $-\Delta u = f(u)$ in $\mathbb{R}^2$, Arch. Rational Mech. Anal., 96 (1986), 147–165.

[8] L. Carleson and A. Sun-Yung Chang, On the existence of an extremal function for an inequality of J. Moser, Bull. Sci. Math., 110(2) (1986), 113–127.

[9] L. Erbe and M. Tang, Uniqueness theorems for positive radial solutions of quasilinear elliptic equations in a ball, J. Differential Equations, 138(2) (1997), 351–379.

[10] B. Gidas, W. M. Ni, and L. Nirenberg, Symmetry and related properties via the maximum principle, Comm. Math. Phys., 68 (1979), 209–243.

[11] J. B. McLeod and L. A. Peletier, Observations on Moser's inequality, Arch. Rational Mech. Anal., 106(3) (1989), 261–285.

[12] Z. Liqun, Uniqueness of positive solutions of $\Delta u + u + u^p = 0$ in a ball, Commun. in Partial Differential Equations, 17(7-8) (1992), 1141-1164.

[13] J. Moser, A sharp form of an inequality by N. Trudinger, Indiana Univ. Math. J., 20 (1970/71), 1077-1092.

[14] E. Nabana, F. de Thelin, On the uniqueness of positive solutions for quasilinear elliptic equations, Nonlinear Anal., 31(3-4) (1998), 413–430.

[15] Wei-Ming Ni and R.D. Nussbaum, Uniqueness and non uniqueness for positive radial solutions of $\Delta u + f(u, r) = 0$, Comm. Pure Appl. Math., 38(1) (1985), 67-108.
[16] P. N. Srikanth, Uniqueness of solutions of nonlinear Dirichlet problems, Differential Integral Equations, 6(3) (1993), 663–670.
[17] M. Tang, Uniqueness of positive radial solutions for n-Laplacian Dirichlet problems, Proc. Roy. Soc. Edinburgh Sect. A, 130(6) (2000), 1405–1416.
[18] H. Volkmer, On the differential equation $y'' + \exp\{y^m - t\} = 0$, Asymptotic Anal., 10(1) (1995), 63–75.

T.I.F.R. CAM, P.B. No. 6503,, Sharadanagar, Chikkabommasandra, Bangalore 560065, India
E-mail address: `adiadimurthi@gmail.com` and `aditi@math.tifrbng.res.in`

Department of Mathematics,, Louisiana State University,, USA
E-mail address: `kadimu1@lsu.edu` and `karthikaditi@gmail.com`

LMAP (UMR CNRS 5142) Bat. IPRA, Avenue de l'Université, F-64013 Pau, France
E-mail address: `jacques.giacomoni@univ-pau.fr`